

TWISTED JETS, MOTIVIC MEASURE AND ORBIFOLD COHOMOLOGY

TAKEHIKO YASUDA

ABSTRACT. We introduce the notion of twisted jets. For a Deligne-Mumford stack $\mathcal{X}$ of finite type over $\mathbb{C}$, a twisted ∞ -jet on $\mathcal{X}$ is a representable morphism $\mathcal{D} \rightarrow \mathcal{X}$ such that $\mathcal{D}$ is a smooth Deligne-Mumford stack with the coarse moduli space $\mathrm{Spec} \mathbb{C}[[t]]$. We study the motivic measure on the space of the twisted ∞ -jets on a smooth Deligne-Mumford stack.

As an application, we prove that two birational minimal models with Gorenstein quotient singularities have the same orbifold cohomology with Hodge structure.

1. INTRODUCTION

In 1995, Kontsevich produced the theory named *motivic integration* [Kon95]. Since then, this remarkable idea has become a powerful method for examining both the local and global structures of varieties.

Let X be a variety over $\mathbb{C}$. For $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, an n -jet on X is a $\mathbb{C}[[t]]/(t^{n+1})$ -point of X , where we have followed the convention $(t^\infty) = (0)$. The n -jets of X naturally constitute a variety (or pro-variety if $n = \infty$), denoted $L_n X$. For $m \geq n$, the natural surjection $\mathbb{C}[[t]]/(t^{m+1}) \rightarrow \mathbb{C}[[t]]/(t^{n+1})$ induces the truncation morphism $L_m X \rightarrow L_n X$.

Consider the case where X is smooth and of dimension d . Then $L_n X$ is a locally trivial affine space bundle over X . (Whenever X is singular, it fails. For example, for $n = 1$, $L_1 X$ is the tangent space of X and hence not a locally trivial bundle over X .) The idea of Kontsevich is to give $L_\infty X$ a measure which takes values in the Grothendieck ring $\mathbb{M}$ of k -varieties which is localized by the class $\mathbb{L}$ of the affine line. For each $n \in \mathbb{Z}_{\geq 0}$, the family of constructible subsets of $L_n X$ is stable under finite union or finite intersection. In other words, this family is a Boolean algebra. The map

$$\begin{aligned} \{\text{constructible subsets of } L_n X\} &\rightarrow \mathbb{M} \\ A &\mapsto \{A\} \mathbb{L}^{-nd} \end{aligned}$$

is a finite additive measure. For each $m > n \in \mathbb{Z}_{\geq 0}$, because the truncation morphism $\pi_n^m : L_m X \rightarrow L_n X$ is a locally trivial affine space bundle of

1991 *Mathematics Subject Classification*. Primary 14F43; Secondary 14B05, 14B10, 14E15.

Key words and phrases. Twisted jets, Deligne-Mumford stacks, motivic measure, orbifold cohomology, Gorenstein quotient singularities.

relative dimension $(m - n)d$, the pull-back

$$(\pi_n^m)^{-1} : \{\text{constructible subsets of } L_n X\} \rightarrow \{\text{constructible subsets of } L_m X\},$$

is considered to be an extension of the measure into a bigger Boolean algebra. The *motivic measure* on $L_\infty X$ is defined to be the limit of these extensions. Denef and Loeser generalized the motivic measure to the case where X is singular [DL99].

The integral of a function with respect to the motivic measure produces a new invariant. In particular, when X is smooth and the function is constant equal to 1, then the integral, which is the full volume of $L_\infty X$, is the class of X in $\mathbb{M}$. It reduces to the Hodge structure of the cohomology of X if X is complete. By the transformation rule of Kontsevich, for any resolution $Z \rightarrow X$, it equals the integral of a function on $L_\infty Z$ determined by the relative canonical divisor $K_{Z/X}$. This implies the following theorem of Kontsevich, which we will generalize:

Theorem 1.1. *Let X and X' be smooth complete varieties. Suppose that there are proper birational morphisms $Z \rightarrow X$ and $Z \rightarrow X'$ such that $K_{Z/X} = K_{Z/X'}$. Then the rational cohomologies of X and X' have the same Hodge structures.*

It is key that by the valuative criterion for properness, almost every ∞ -jet on X lifts to a unique ∞ -jet on Z , and hence the map $L_\infty Z \rightarrow L_\infty X$ is bijective outside of measure zero subsets. Note that Batyrev first proved the equality of Betti numbers in the case where X and X' are Calabi-Yau varieties, with p -adic integration and the Weil conjecture [Bat99a].

Let X be a variety with Gorenstein canonical singularities. Denef and Loeser gave $L_\infty X$ another measure, called the *motivic Gorenstein measure*, denoted μ_X^{Gor} [DL02]. As in the case with X smooth, $\mu_X^{\text{Gor}}(L_\infty X)$ is calculated by the relative canonical divisor $K_{Z/X}$ for a resolution $Z \rightarrow X$. This implies:

Proposition 1.2. *Let X and X' be varieties with Gorenstein canonical singularities. Suppose that there are proper birational morphisms $Z \rightarrow X$ and $Z \rightarrow X'$ such that $K_{Z/X} = K_{Z/X'}$. Then $\mu_X^{\text{Gor}}(L_\infty X) = \mu_{X'}^{\text{Gor}}(L_\infty X')$*

Quotient singularities form one of the mildest classes of singularities¹. If X is a variety with quotient singularities, then we can give X an orbifold structure. In the algebro-geometric context, there is a smooth Deligne-Mumford stack $\mathcal{X}$ such that X is the coarse moduli space of $\mathcal{X}$ and the automorphism group of general points of $\mathcal{X}$ is trivial. Although the natural morphism $\mathcal{X} \rightarrow X$ is proper and birational, *not* almost every ∞ -jet on X lifts to a $\mathbb{C}[[t]]$ -point of $\mathcal{X}$ from lack of the *strict* valuative criterion for properness. However by twisting the source $\text{Spec } \mathbb{C}[[t]]$, we can lift almost every ∞ -jet on X to $\mathcal{X}$. More precisely, A *twisted* ∞ -jet on $\mathcal{X}$ is a representable morphism

¹Here the words ‘quotient singularities’ mean ‘quotient singularities with respect to the étale topology’, see Definition 4.27.

$\mathcal{D} \rightarrow \mathcal{X}$ such that $\mathcal{D}$ is a smooth Deligne-Mumford stack with the coarse moduli space $\mathrm{Spec} \mathbb{C}[[t]]$ and $\mathcal{D}$ contains $\mathrm{Spec} \mathbb{C}((t))$ as open substack. (A paper of Abramovich and Vistoli [AV02] was the inspiration for this notion. They introduced the notion of twisted stable map.) For almost every ∞ -jet $\gamma : \mathrm{Spec} \mathbb{C}[[t]] \rightarrow X$, there is a unique twisted ∞ -jet $\mathcal{D} \rightarrow \mathcal{X}$ such that the induced morphism $\mathrm{Spec} \mathbb{C}[[t]] \rightarrow X$ of the coarse moduli spaces is γ . If $\overline{\mathcal{L}_\infty \mathcal{X}}$ is the coarse moduli space of the twisted ∞ -jets on $\mathcal{X}$, then we define the *motivic measure* $\mu_{\mathcal{X}}$ on $\overline{\mathcal{L}_\infty \mathcal{X}}$ in a similar fashion as on $L_\infty X$, though it takes values in the Grothendieck ring of Hodge structures. We show the following close relation between μ_X^{Gor} and $\mu_{\mathcal{X}}$:

Theorem 1.3. *The following equality holds:*

$$\mu_X^{\mathrm{Gor}} = \sum_{\mathcal{Y} \subset I(\mathcal{X})} \mathbb{L}^{s(\mathcal{Y})} \mu_{\mathcal{X}}.$$

For the precise meaning, see Theorem 3.15.

Chen and Ruan defined the *orbifold cohomology* for arbitrary orbifold [CR00]. It originates from string theory on orbifolds [DHVW]. Let X be a variety with Gorenstein quotient singularities and $\mathcal{X}$ as above. The *inertia stack* of $\mathcal{X}$, denoted $I(\mathcal{X})$, is an object in the algebro-geometric realm that corresponds to the twisted sector. We define the i -th orbifold cohomology group of X by:

$$H_{\mathrm{orb}}^i(X, \mathbb{Q}) := \bigoplus_{\mathcal{Y} \subset I(\mathcal{X})} H^{i-2s(\mathcal{Y})}(\overline{\mathcal{Y}}, \mathbb{Q}) \otimes \mathbb{Q}(-s(\mathcal{Y})),$$

where $\mathcal{Y}$ runs over the connected components of $I(\mathcal{X})$, $\overline{\mathcal{Y}}$ is the coarse moduli space of $\mathcal{Y}$ and $s(\mathcal{Y})$ is an integer which is representation-theoretically determined.

- Remark 1.4.* (1) The author guesses, though is not sure, that our orbifold cohomology is equal to one by Chen and Ruan in [CR00]. The point he wonders is whether the cohomology groups of $\overline{\mathcal{Y}}$ are isomorphic to ones of the analytic orbifold (V-manifold) associated to $\mathcal{Y}$.
- (2) The orbifold Hodge numbers (that is, the Hodge numbers associated to $H_{\mathrm{orb}}^i(X, \mathbb{Q})$) are equal to Batyrev's stringy Hodge numbers [Bat99b]. It follows from Lemma 2.16 and Theorem 1.3.

Theorem 1.3 implies the fact that when X is complete, the invariant $\mu_X^{\mathrm{Gor}}(L_\infty X)$ reduces to the alternating sum of the orbifold cohomology groups of X . Hence we obtain the following, conjectured by Ruan [Rua00]:

Theorem 1.5 (=Corollary 3.16). *Let X and X' be complete varieties with Gorenstein quotient singularities. Suppose that there are proper birational morphisms $Z \rightarrow X$ and $Z \rightarrow X'$ such that $K_{Z/X} = K_{Z/X'}$. Then the orbifold cohomologies of X and X' have the same Hodge structures.*

If X and X' are birational minimal models, that is, K_X and K'_X are nef, then for a common resolution Z of X and X' , we have the equality $K_{Z/X} = K_{Z/X'}$ (see [KM98, Prop. 3.51]). Hence X and X' have the same orbifold cohomology with Hodge structure. Note that in the case where X and X' are global quotients, Theorem 1.5 is due to Batyrev [Bat99b] and Denef-Loeser [DL02]. After writing out the first version of this paper, I learned by an e-mail message from Ernesto Lupercio that Mainak Poddar and he independently proved Theorem 1.5.

Contents. The paper is organized as follows. In Section 1, we review motivic measures. Section 2 is the central part of the paper. Here we introduce the notion of a twisted jet and examine the space of them. Then we prove the main result. In Section 3, we review Deligne-Mumford stacks and prove some general results on Deligne-Mumford stacks which we need in the preceding section.

Conventions and Notations.

- In Section 1 and 2, we work over $\mathbb{C}$.
- For a Deligne-Mumford stack $\mathcal{X}$, we denote by $\overline{\mathcal{X}}$ the coarse moduli space of $\mathcal{X}$.
- We denote by (Sch/S) (resp. $(\text{Sch}/\mathbb{C})$) the category of schemes over a scheme S (resp. over $\mathbb{C}$).
- For a $\mathbb{C}$ -scheme X (or more generally a stack over $\mathbb{C}$) and a $\mathbb{C}$ -algebra R , we denote by $X \otimes R$ the product $X \times_{\mathbb{C}} \text{Spec } R$. Then we denote by $X[[t]]$ (resp. $X[[t^{1/l}]]$) the scheme $X \otimes \mathbb{C}[[t]]$ (resp. $X \otimes \mathbb{C}[[t^{1/l}]]$).

Acknowledgements. This paper forms a part of my master's thesis. I am deeply grateful to Yujiro Kawamata for recommending me to learn this area and for his direction and encouragement. I also would like to thank James McKernan and Hokuto Uehara for reading this paper and pointing out English mistakes.

2. MOTIVIC MEASURE; REVIEW

In this section, we would like to review the theory of motivic measures, developed by Kontsevich [Kon95], Batyrev [Bat99b], Denef and Loeser [DL99], [DL02]. I mention [Cra99] for a nice introduction and [Loo02], [DL01] for surveys.

2.1. Completing Grothendieck rings. Let us first construct the ring in which motivic measures take values.

A variety means a reduced scheme of finite type over $\mathbb{C}$.

Definition 2.1. We define the *Grothendieck ring of varieties*, denoted $K_0(\text{Var})$, to be the abelian group generated by the isomorphism classes $\{X\}$ of varieties with the relations $\{X\} = \{X \setminus Y\} + \{Y\}$ if Y is a closed subvariety of X . The ring structure is defined by $\{X\}\{Y\} = \{X \times Y\}$.

In the same fashion, we can define the Grothendieck ring of separated algebraic spaces of finite type. Actually, it is the same thing as $K_0(\text{Var})$, since every noetherian algebraic space decomposes into the disjoint union of schemes [Knu71, Prop. 6.6].

Suppose that A is a constructible subset of a variety X , that is, A is a disjoint union of locally closed subvarieties $A_i \subset X$. Then we put $\{A\} := \sum_i \{A_i\} \in K_0(\text{Var})$, which is independent of the choice of stratifications. We denote the class of $\mathbb{A}^1$ by $\mathbb{L}$ and the localization $K_0(\text{Var})[\mathbb{L}^{-1}]$ by $\mathbb{M}$. For $m \in \mathbb{Z}$, let $F_m \mathbb{M}$ be the subgroup of $\mathbb{M}$ generated by the elements $\{X\} \mathbb{L}^{-i}$ with $\dim X - i \leq -m$. The collection $(F_m \mathbb{M})_{m \in \mathbb{Z}}$ is a descending filtration of $\mathbb{M}$ with

$$(2.1) \quad F_m \mathbb{M} \cdot F_n \mathbb{M} \subset F_{m+n} \mathbb{M}.$$

Definition 2.2. We define the *complete Grothendieck ring of varieties* by

$$\hat{\mathbb{M}} := \varprojlim \mathbb{M} / F_m \mathbb{M}.$$

By condition (2.1), it has a natural ring structure.

Note that it is not known whether the natural map $\mathbb{M} \rightarrow \hat{\mathbb{M}}$ is injective or not.

Recall that a *Hodge structure* is a finite dimensional $\mathbb{Q}$ -vector space H with a bigrading $H \otimes \mathbb{C} = \bigoplus_{p,q \in \mathbb{Z}} H^{p,q}$ such that $H^{p,q}$ is the complex conjugate of $H^{q,p}$ and each weight summand $\bigoplus_{p+q=m} H^{p,q}$ is defined over $\mathbb{Q}$. The category HS of Hodge structures is an abelian category with tensor product.

Definition 2.3. We define the *Grothendieck ring of Hodge structures*, denoted $K_0(\text{HS})$, to be the abelian group which consists of formal differences $\{H\} - \{H'\}$, where $\{H\}$ and $\{H'\}$ are isomorphism classes of Hodge structures. The addition and the multiplication come from $\oplus$ and $\otimes$ respectively.

A *mixed Hodge structure* is a finite dimensional $\mathbb{Q}$ -vector space H with increasing filtration $W_\bullet H$, called the *weight filtration*, such that the associated graded $\text{Gr}_\bullet^W H$ underlies a Hodge structure having $\text{Gr}_m^W H$ as weight m summand. For a mixed Hodge structure H , we denote by $\{H\}$ the element $\{\text{Gr}_\bullet^W H\}$ of $K_0(\text{HS})$.

The cohomology groups $H_c^i(X, \mathbb{Q})$ with compact supports of a variety X has a natural mixed Hodge structure.

Definition 2.4. We define the *Hodge characteristic* $\chi_h(X)$ of X by

$$\chi_h(X) := \sum_i (-1)^i \{H_c^i(X, \mathbb{Q})\} \in K_0(\text{HS}).$$

Consider the following map:

$$(\text{Varieties}) \rightarrow K_0(\text{HS}), X \mapsto \chi_h(X).$$

It factors through the map

$$(\text{Varieties}) \rightarrow \mathbb{M}, X \mapsto \{X\},$$

because the following hold:

- $\chi_h(X \times Y) = \chi_h(X)\chi_h(Y)$,
- $\chi_h(X) = \chi_h(X \setminus Y) + \chi_h(Y)$ if $Y \subset X$ is closed,
- the Hodge characteristic of the affine line, $\chi_h(\mathbb{A}^1) = \{H_c^2(\mathbb{A}^1, \mathbb{Q})\}$, is invertible.

We also denote by χ_h the induced homomorphism $\mathbb{M} \rightarrow K_0(\text{HS})$.

For $m \in \mathbb{Z}$, let $F_m K_0(\text{HS})$ be the subgroup generated by the elements $\{H\}$ such that the maximum weight of H is less than or equal to $-m$.

Definition 2.5. We define the *complete Grothendieck ring of Hodge structures*, denoted $K_0(\hat{\text{HS}})$, as follows:

$$K_0(\hat{\text{HS}}) := \varprojlim K_0(\text{HS})/F_m K_0(\text{HS}).$$

We can see that the natural map $K_0(\text{HS}) \rightarrow K_0(\hat{\text{HS}})$ is injective. Because the maximal weight of $H^i(X, \mathbb{Q})$ does not exceed $2 \dim X$, χ_h extends to $\chi_h : \hat{\mathbb{M}} \rightarrow K_0(\hat{\text{HS}})$.

2.2. Jets on schemes. For convenience's sake, we denote by (t^∞) the ideal (0) of the power series ring $\mathbb{C}[[t]]$. For $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, we denote by D_n the affine scheme $\text{Spec } \mathbb{C}[[t]]/(t^{n+1})$.

Definition 2.6. Let X be a scheme. For $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, we define the *scheme of n -jets*² of X , denoted $L_n X$, to be the scheme representing the functor

$$\begin{aligned} (\text{Sch}/\mathbb{C}) &\rightarrow (\text{Sets}) \\ U &\mapsto \text{Hom}_{(\text{Sch}/\mathbb{C})}(U \times D_n, X). \end{aligned}$$

Greenberg [Gre61] proved the representability of the functor for $n < \infty$. For $m, n \in \mathbb{Z}_{\geq 0}$ with $m < n$, a canonical closed immersion $D_m \rightarrow D_n$ induces a canonical projection $L_n X \rightarrow L_m X$. Since all these projections are affine morphisms, the projective limit $L_\infty X = \varprojlim L_n X$ exists in the category of schemes.

If X is of finite type, then for $n < \infty$, so is $L_n X$. If X is smooth and of pure dimension d , then, for each $n \in \mathbb{Z}_{\geq 0}$, the natural projection $L_{n+1} X \rightarrow L_n X$ is a Zariski locally trivial $\mathbb{A}^d$ -bundle. If $f : Y \rightarrow X$ is a morphism of schemes, then for each n , there is a canonical morphism $f_n : L_n Y \rightarrow L_n X$.

2.3. Motivic measure. Let X be a scheme of pure dimension d . By abuse of notation, we denote the set of points of $L_\infty X$ also by $L_\infty X$. Let $\pi_n : L_\infty X \rightarrow L_n X$ be the canonical projection.

Definition 2.7. A subset A of $L_\infty X$ is *stable at level n* if we have:

- (1) $\pi_n(A)$ is a constructible subset in $L_n X$,

²In [DL99], an ∞ -jet is called an *arc*. As it is more convenient, I prefer my terminology.

- (2) $A = \pi_n^{-1}\pi_n(A)$,
- (3) for any $m \geq n$, the projection $\pi_{m+1}(A) \rightarrow \pi_m(A)$ is a piecewise trivial $\mathbb{A}^d$ -bundle.

(A morphism $f : Y \rightarrow X$ of schemes is called a *piecewise trivial $\mathbb{A}^d$ -bundle* if there is a stratification $X = \coprod X_i$ such that $f|_{f^{-1}(X_i)} : f^{-1}(X_i) \rightarrow X_i$ is isomorphic to $X_i \times \mathbb{A}^d \rightarrow X_i$ for each i .) A subset A of $L_\infty X$ is *stable* if it is stable at level n for some $n \in \mathbb{Z}_{\geq 0}$.

The stable subsets of $L_\infty X$ constitute a Boolean algebra. If $A \subset L_\infty X$ is a stable subset, then $\{\pi_m(A)\}\mathbb{L}^{-md} \in \hat{\mathbb{M}}$ is constant for $m \gg 0$. We denote it by $\mu_X(A)$ ³. The map

$$\begin{aligned} \mu_X : \{\text{stable subsets of } L_\infty X\} &\rightarrow \hat{\mathbb{M}} \\ A &\mapsto \mu_X(A) \end{aligned}$$

is a finite additive measure. Let us extend this measure to a bigger family of subsets of $L_\infty X$.

Definition 2.8. A subset $A \subset L_\infty X$ is called *measurable* if, for every $m \in \mathbb{Z}$, there are stable subsets $A_m \subset L_\infty X$ and $C_i \subset L_\infty X$, $i \in \mathbb{Z}_{>0}$ such that the symmetric difference $(A \cup A_m) \setminus (A \cap A_m)$ is contained in $\cup_i C_i$ and we have $\mu_X(C_i) \in F_m \hat{\mathbb{M}}$ for all i , and $\lim_{i \rightarrow \infty} \mu_X(C_i) = 0$ in $\hat{\mathbb{M}}$.

The measurable subsets of $L_\infty X$ also constitute a Boolean algebra. Suppose that $A \subset L_\infty X$ is measurable and $A_m \subset L_\infty X$, $m \in \mathbb{Z}$ are stable subsets as in the definition above. We put $\mu_X(A) := \lim_{m \rightarrow \infty} \mu_X(A_m)$. It is independent of the choice of A_m , see [Loo02, Prop. 2.2], [DL02, Th. A.6]. The map

$$\begin{aligned} \mu_X : \{\text{measurable subsets of } L_\infty X\} &\rightarrow \hat{\mathbb{M}} \\ A &\mapsto \mu_X(A) \end{aligned}$$

is a finite additive measure.

Definition 2.9. We call this the *motivic measure* on $L_\infty X$.

Definition 2.10. Let $A \subset L_\infty X$ be a measurable subset and $\nu : A \rightarrow \mathbb{Z} \cup \{\infty\}$ a function. We say that ν is a *measurable function* if the fibers are measurable and $\mu_X(\nu^{-1}(\infty)) = 0$. For a measurable function ν , we formally define the *motivic integration* of $\mathbb{L}^\nu$ by

$$\int_A \mathbb{L}^\nu d\mu_X := \sum_{n \in \mathbb{Z}} \mu_X(\nu^{-1}(n)) \mathbb{L}^n.$$

We say that $\mathbb{L}^\nu$ is *integrable* if the series above converges in $\hat{\mathbb{M}}$.

³This differs from the definition in [Cra99], [Bat99b] and [DL99] by a factor $\mathbb{L}^d$.

Example 2.11. Let $\mathcal{J}$ be an ideal sheaf on X . A point $\gamma \in L_\infty X$ corresponds to a morphism $\text{Spec } \kappa \rightarrow L_\infty X$ for the residue field κ of γ and hence to a morphism $\gamma' : \text{Spec } \kappa[[t]] \rightarrow X$. The function

$$\begin{aligned} \text{ord } \mathcal{J} : L_\infty X &\rightarrow \mathbb{Z}_{\geq 0} \cup \{\infty\} \\ \gamma &\mapsto n \text{ if } (\gamma')^{-1}\mathcal{J} = (t^n) \end{aligned}$$

is a measurable function by the following lemma.

Lemma 2.12. [Loo02, Prop. 3.1], [DL99, Lem. 4.4]. *For a subvariety $Y \subset X$ of positive codimension, the subset $L_\infty Y \subset L_\infty X$ is of measure zero.*

Example 2.13. [Bat99b, Th. 3.6], [Cra99, Th. 2.15]. Let X be a smooth variety of dimension d and $E = \sum_{i=1}^r d_i E_i$ an effective divisor on X with simple normal crossing support. For a subset $J \subset \{1, \dots, r\}$, we define

$$E_J^\circ := \bigcap_{i \in J} E_i \setminus \bigcup_{i \in \{1, \dots, r\} \setminus J} E_i.$$

If $\mathcal{J}_E$ is the ideal sheaf associated to E , then we have the following formula:

$$\int_{L_\infty X} \mathbb{L}^{-\text{ord } \mathcal{J}_E} d\mu_X = \sum_{J \subset \{1, \dots, r\}} \{E_J^\circ\} \prod_{i \in J} \frac{\mathbb{L} - 1}{\mathbb{L}^{d_i+1} - 1}.$$

2.4. The transformation rule. Let X and Y be varieties of pure dimension d and $f : Y[[t]] \rightarrow X[[t]]$ a morphism over D_∞ . We define the morphisms $f_n : L_n Y \rightarrow L_n X$ as follows. For a scheme U , a U -point γ of $L_n Y$ is a morphism $\gamma' : D_n \times U \rightarrow Y$. If ψ is the composition of natural morphisms $D_n \times U \xrightarrow{\text{pr}} D_n \rightarrow D_\infty$, then we define $f_n(\gamma)$ to be the U -point of $L_n X$ corresponding to the composition

$$D_n \times U \xrightarrow{(\gamma', \psi)} Y \times D_\infty = Y[[t]] \xrightarrow{f} X[[t]] \rightarrow X.$$

Assume that Y is smooth. We define the *jacobian ideal sheaf* $\mathcal{J}_f$ of f to be the 0-th Fitting ideal sheaf of $\Omega_{Y[[t]]/X[[t]]}$ (see [EH99, V.1.3] or [Eis95, 20.2]). This is the ideal sheaf such that $\mathcal{J}_f \Omega_{Y[[t]]/D_\infty}^d = f^*(\Omega_{X[[t]]/D_\infty}^d / (\text{tors}))$ as subsheaves of $\Omega_{Y[[t]]/D_\infty}^d$, where (tors) is the torsion. The following, called the *transformation rule*, is the most basic theorem in the theory.

Theorem 2.14. [DL02, Th. 1.16], [Loo02, Th. 3.2]. *Let A be a measurable set in $L_\infty Y$ and $\nu : f_\infty(A) \rightarrow \mathbb{Z} \cup \{\infty\}$ a measurable function. Suppose that $f_\infty|_A$ is injective. Then we have the following equality:*

$$\int_{f_\infty(A)} \mathbb{L}^\nu d\mu_X = \int_A \mathbb{L}^{\nu \circ f_\infty - \text{ord } \mathcal{J}_f} d\mu_Y.$$

We will generalize this later (Theorem 3.18).

2.5. The motivic Gorenstein measure. Let X be a variety with 1-Gorenstein and canonical singularities, that is, the canonical sheaf ω_X is invertible and all discrepancies are ≥ 0 , (see [KMM87, §0-2]). Then there exists a natural morphism $\Omega_X^d \rightarrow \omega_X$. The kernel of this morphism is the torsion. We define an ideal sheaf $\mathcal{J}_X$ on X by the equation:

$$\mathcal{J}_X \omega_X = \text{Im}(\Omega_X^d \rightarrow \omega_X).$$

Then $\mathbb{L}^{\text{ord } \mathcal{J}_X}$ is integrable by Example 2.13 and Lemma 2.16.

Definition 2.15. We define the *motivic Gorenstein measure* μ_X^{Gor} on $L_\infty X$ as follows:

$$\begin{aligned} \mu_X^{\text{Gor}} : \{\text{measurable subsets of } L_\infty X\} &\rightarrow \mathbb{M} \\ A &\mapsto \int_A \mathbb{L}^{\text{ord } \mathcal{J}_X} d\mu_X. \end{aligned}$$

Lemma 2.16. Let X and X' be complete varieties with 1-Gorenstein canonical singularities.

- (1) Let A be a measurable subset of $L_\infty X$ and $f : Z \rightarrow X$ be a resolution. Then

$$\int_A \mathbb{L}^{\text{ord } \mathcal{J}_X} d\mu_X = \int_{f_\infty^{-1}(A)} \mathbb{L}^{-\text{ord } \mathcal{K}} d\mu_Z,$$

where $\mathcal{K}$ is the ideal sheaf associated to $K_{Z/X}$.

- (2) Suppose that there exist proper birational morphisms $f : Z \rightarrow X$ and $g : Z \rightarrow X'$ with $K_{Z/X} = K_{Z/X'}$. Then we have $\mu_X^{\text{Gor}}(L_\infty X) = \mu_{X'}^{\text{Gor}}(L_\infty X')$.

Proof. (1). By Theorem 2.14, we have

$$\int_A \mathbb{L}^{\text{ord } \mathcal{J}_X} d\mu_X = \int_{f_\infty^{-1}(A)} \mathbb{L}^{\text{ord } \mathcal{J}_X \circ f_\infty - \text{ord } \mathcal{J}_f} d\mu_Z.$$

We have to show

$$(2.2) \quad \text{ord } \mathcal{J}_X \circ f_\infty - \text{ord } \mathcal{J}_f = -\text{ord } \mathcal{K}.$$

Pulling back $\mathcal{J}_X \omega_X \cong \Omega_X^d / (\text{tors})$, we have $(f^{-1} \mathcal{J}_X)(f^* \omega_X) \cong \mathcal{J}_f \Omega_Z^d$. On the other hand, we have $f^* \omega_X \cong \mathcal{K} \omega_Z \cong \mathcal{K} \Omega_Z^d$. Hence $(f^{-1} \mathcal{J}_X) \cdot \mathcal{K} = \mathcal{J}_f$. This shows (2.2).

(2) is a direct consequence of (1). $\square$

Remark 2.17. The invariant $\mu_X^{\text{Gor}}(L_\infty X)$ has been already introduced in [Kon95] by using a resolution of singularities. Denef and Loeser constructed the invariant more directly with the motivic Gorenstein measure [DL02].

Suppose X is complete. The question is whether $\chi_h \circ \mu_X^{\text{Gor}}(L_\infty X)$ is the alternating sum of a *kind of cohomology groups*, as the case where X is smooth. It is known that when X is a global quotient, the answer is Yes [Bat99b], [DL02]. Our result, Theorem 3.15, says that when X has only quotient singularities, the answer is Yes.

3. TWISTED JETS

In this section, we need manage the theory of Deligne-Mumford stacks. See the next section for the generalities about Deligne-Mumford stack.

3.1. Non-twisted jets on stacks. The following is a direct generalization of the notion of jet on schemes:

Definition 3.1. Let $\mathcal{X}$ be a Deligne-Mumford stack. For $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, we define the *stack of non-twisted n -jets* of $\mathcal{X}$, denoted $L_n\mathcal{X}$, as follows. An object of $L_n\mathcal{X}$ over $U \in (\text{Sch}/\mathbb{C})$ is an object of $\mathcal{X}$ over $U \times D_n$. For a morphism $\varphi : V \rightarrow U$ in $(\text{Sch}/\mathbb{C})$, a morphism in $L_n\mathcal{X}$ over φ is a morphism in $\mathcal{X}$ over $\varphi \times \text{id}_{D_n}$.

Lemma 3.2. *For every $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, $L_n\mathcal{X}$ is a stack.*

Proof. It is clear that they satisfy the axioms of category fibered in groupoids. If $(U_i \rightarrow U)_i$ is an étale covering in $(\text{Sch}/\mathbb{C})$, then so is $(U_i \times D_n \rightarrow U \times D_n)_i$. As a result, they satisfy the axioms of stack, too. $\square$

Let $f : Y \rightarrow X$ be a morphism of schemes and $Z \subset Y$ a closed subscheme with an ideal sheaf $\mathfrak{a}$. We say that f is *Z -étale* if for any ring A and any nilpotent ideal $J \subset A$, and for any commutative diagram of solid arrows

$$\begin{array}{ccc} \text{Spec } A/J & \xrightarrow{\varphi} & Y \\ \downarrow & \nearrow & \downarrow \\ \text{Spec } A & \longrightarrow & X \end{array}$$

such that $\varphi^{-1}\mathfrak{a}$ is nilpotent, there is a unique broken arrow which makes the whole diagram commutative. Z -étaleness is defined for a representable morphism of stacks also in the evident fashion.

Lemma 3.3. (1) *Let M be a scheme and $N \subset M$ a closed subscheme. We denote by $(L_nM)_N$ the subscheme of L_nM parametrizing the jets with the base point in N . Let $p : M \rightarrow \mathcal{X}$ be an N -étale morphism. Then, for every $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, we have a natural isomorphism:*

$$(L_nM)_N \cong L_n\mathcal{X} \times_{\mathcal{X}} N.$$

(2) *For every $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, $L_n\mathcal{X}$ is a Deligne-Mumford stack.*

Proof. (1). Let us first show $L_n\mathcal{X} \times_{\mathcal{X}} N$ is an algebraic space. Let $\pi : L_n\mathcal{X} \rightarrow \mathcal{X}$ be the canonical projection. An object of $L_n\mathcal{X} \times_{\mathcal{X}} N$ is a triple (γ, f, α) where $\gamma : U \times D_n \rightarrow \mathcal{X}$, $f : U \rightarrow N$, and $\alpha : \pi(\gamma) \rightarrow p(f)$ is a morphism in $\mathcal{X}$ over U . By definition, an automorphism of (γ, f, α) is an automorphism θ of γ such that $\alpha \circ \pi(\theta) = p(\text{id}_f) \circ \alpha$. Hence θ must be the identity. We have thus proved that the automorphism of every object of $L_n\mathcal{X} \times_{\mathcal{X}} N$ is trivial and hence that $L_n\mathcal{X} \times_{\mathcal{X}} N$ is an algebraic space (see [LMB00, Cor. 8.1.1]).

The diagram of solid arrows

$$\begin{array}{ccccc}
 U & \xrightarrow{f} & N & \hookrightarrow & M \\
 \downarrow & & \searrow \tau & & \downarrow p \\
 U \times D_n & \xrightarrow{\gamma} & & & \mathcal{X}
 \end{array}$$

is commutative. We can see that there is a unique broken arrow τ in the diagram. If $n < \infty$, since p is N -étale, this is trivial from the definition. If $n = \infty$, since D_∞ is the direct limit of D_n , $0 \leq n < \infty$, this follows from the case $n < \infty$. Sending (γ, f, α) to τ defines a morphism $\Phi : L_n \mathcal{X} \times_{\mathcal{X}} M \rightarrow L_n M$.

The inverse of the morphism Φ is given by $(p_n, \pi_M) : (L_n M)_N \rightarrow L_n \mathcal{X} \times_{\mathcal{X}} N$ where $p_n : L_n M \rightarrow L_n \mathcal{X}$ and $\pi_M : (L_n M)_N \rightarrow N = (L_0 M)_N$ are the natural morphisms. We have thus proved (1).

(2). Now suppose that p is étale and surjective. Consider the following cartesian diagram:

$$\begin{array}{ccc}
 L_n M & \longrightarrow & L_n \mathcal{X} \\
 \pi_M \downarrow & \square & \downarrow \pi \\
 M & \xrightarrow{p} & \mathcal{X}.
 \end{array}$$

Because π_M is representable and p is étale and surjective, π is representable (see [LMB00, Lem. 4.3.3]). This completes the proof (see [LMB00, Prop. 4.5]). $\square$

3.2. Twisted jets. For a positive integer l , we put $\zeta_l := \exp(2\pi\sqrt{-1}/l)$. Let $\mu_l = \langle \zeta_l \rangle$ be the group of the l -th roots of 1. μ_l acts on D_n by $\zeta_l : t \mapsto \zeta_l t$. We denote by $\mathcal{D}_n^l$ the quotient stack $[D_n/\mu_l]$ with $m = nl$. The stack $\mathcal{D}_n^l$ has the canonical atlas $D_n \rightarrow \mathcal{D}_n^l$ and the closed point $\text{Spec } \mathbb{C} \rightarrow \mathcal{D}_n^l$. We fix a morphism $\mathcal{D}_n^l \rightarrow D_n$ such that D_n is the coarse moduli space of $\mathcal{D}_n^l$ for this morphism, and such that the composition

$$D_n \rightarrow \mathcal{D}_n^l \rightarrow D_n$$

is given by the ring homomorphism $\mathbb{C}[[t]]/(t^{n+1}) \rightarrow \mathbb{C}[[t]]/(t^{m+1})$, $t \mapsto t^l$.

Definition 3.4. Let $\mathcal{X}$ be a Deligne-Mumford stack. A *twisted n -jet of order l* on $\mathcal{X}$ is a representable morphism $\mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$ for an algebraically closed field $\Omega \supset \mathbb{C}$.

For a Deligne-Mumford stack $\mathcal{X}$, the *inertia stack* of $\mathcal{X}$, denoted $I(\mathcal{X})$, is the stack parametrizing the pairs (ξ, α) such that $\xi \in \text{ob } \mathcal{X}$ and $\alpha \in \text{Aut}(\xi)$. For details on inertia stack, see Subsection 4.3. There is a natural forgetting morphism $I(\mathcal{X}) \rightarrow \mathcal{X}$. For $l \in \mathbb{Z}_{>0}$, let $I^l(\mathcal{X}) \subset I(\mathcal{X})$ denote the substack parametrizing the pairs (ξ, α) with $\text{ord}(\alpha) = l$.

Let $\gamma : \mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$ be a twisted n -jet of order l on $\mathcal{X}$. The canonical morphism

$$\tilde{\gamma} : D_n \otimes \Omega \rightarrow \mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$$

is considered to be an Ω -point of $L_m\mathcal{X}$ and the canonical morphism

$$\bar{\gamma} : \text{Spec } \Omega \rightarrow D_m \otimes \Omega \rightarrow \mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$$

to be an Ω -point of $\mathcal{X}$. Since the automorphism group of the closed point of $\mathcal{D}_n^l \otimes \Omega$ is identified with μ_l , γ induces an injection $\mu_l \rightarrow \text{Aut}(\bar{\gamma})$. If $b \in \text{Aut}(\bar{\gamma})$ is the image of ζ_l , then the pair $(\bar{\gamma}, b)$ is regarded as an Ω -point of $I^l(\mathcal{X})$ and the triple $(\tilde{\gamma}, (\bar{\gamma}, b), \text{id}_{\bar{\gamma}})$ to be an Ω -point of $L_m\mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})$. We define a map Ψ by

$$\begin{aligned} \Psi : \{\text{twisted } n\text{-jets of order } l \text{ on } \mathcal{X}\} &\rightarrow |L_m\mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})| \\ \gamma &\mapsto (\tilde{\gamma}, (\bar{\gamma}, b), \text{id}_{\bar{\gamma}}). \end{aligned}$$

Lemma 3.5. *The subset $\text{Im}(\Psi) \subset |L_m\mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})|$ is closed for the Zariski topology.*

Proof. Fix an atlas $p : M \rightarrow \mathcal{X}$ with M separated.

We will first characterize the points in $\text{Im}(\Psi)$. On account of the arguments on groupoid spaces in Subsection 4.1, we can see that the following are equivalent:

- (1) to give a commutative diagram

$$\begin{array}{ccc} D_m \otimes \Omega & \xrightarrow{\eta} & M \\ \downarrow & & \downarrow p \\ \mathcal{D}_n^l \otimes \Omega & \xrightarrow{\gamma} & \mathcal{X} \end{array}$$

such that γ is a twisted n -jet of order l ,

- (2) to give a morphism of groupoid spaces

$$\begin{array}{ccc} (D_m \otimes \Omega) \times \mu_l & \xrightarrow{\delta'} & M \times_{\mathcal{X}} M \\ \text{pr} \downarrow \downarrow \mu_l\text{-action} & & \text{pr}_1 \downarrow \downarrow \text{pr}_2 \\ D_m \otimes \Omega & \xrightarrow{\eta} & M \end{array}$$

such that the composition

$$\text{Spec } \Omega \hookrightarrow (D_m \otimes \Omega) \times \{\zeta_l\} \hookrightarrow (D_m \otimes \Omega) \times \mu_l \xrightarrow{\delta'} M \times_{\mathcal{X}} M$$

corresponds to an automorphism of order l of the following Ω -point of $\mathcal{X}$:

$$\text{Spec } \Omega \hookrightarrow D_m \otimes \Omega \xrightarrow{\eta} M \xrightarrow{p} \mathcal{X}.$$

- (3) to give a morphism $\delta : D_m \otimes \Omega \rightarrow M \times_{\mathcal{X}} M$ such that $\text{pr}_1 \circ \delta \circ \zeta_l = \text{pr}_2 \circ \delta$ and the composition

$$\text{Spec } \Omega \hookrightarrow D_m \otimes \Omega \xrightarrow{\delta} M \times_{\mathcal{X}} M$$

corresponds to an automorphism of order l of the following Ω -point of $\mathcal{X}$:

$$\text{Spec } \Omega \hookrightarrow D_m \otimes \Omega \xrightarrow{\delta} M \times_{\mathcal{X}} M \xrightarrow{\text{pr}_1} M \xrightarrow{p} \mathcal{X}.$$

Any point of $|L_m \mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})|$ is represented by the triple $(\psi, (\overline{\psi}, b), \text{id}_{\overline{\psi}})$ such that ψ is an Ω -point of $L_m \mathcal{X}$ with an algebraically closed field Ω , $\overline{\psi}$ is an Ω -point of $\mathcal{X}$ corresponding to the composition

$$\text{Spec } \Omega \hookrightarrow D_m \otimes \Omega \xrightarrow{\psi} \mathcal{X}$$

and b is an automorphism of $\overline{\psi}$. Then the equivalence above implies:

♣ $(\psi, (\overline{\psi}, b), \text{id}_{\overline{\psi}})$ is in $\text{Im}(\Psi)$ iff for a lift $\eta : D_m \otimes \Omega \rightarrow M$ of ψ , there exists a morphism $\delta : D_m \otimes \Omega \rightarrow M \times_{\mathcal{X}} M$ such that $\text{pr}_1 \circ \delta = \eta$ and $\text{pr}_2 \circ \delta = \eta \circ \zeta_l$ and the composition $\text{Spec } \Omega \rightarrow D_m \otimes \Omega \xrightarrow{\delta} M \times_{\mathcal{X}} M$ corresponds to b .

Let ξ and σ be points of $L_m \mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})$. Suppose that σ is in $\text{Im}(\Psi)$ and ξ is a specialization of σ . It suffices to show that ξ is in $\text{Im}(\Psi)$. By [LMB00, Prop. 7.2.1], there is a complete discrete valuation ring R with algebraically closed residue field κ and quotient field K such that there is a commutative diagram as follows:

$$\begin{array}{ccc} \text{Spec } K & & \\ \downarrow & \searrow \sigma & \\ \text{Spec } R & \xrightarrow{\theta} & L_m \mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X}) \\ \uparrow & \nearrow \xi & \\ \text{Spec } \kappa & & \end{array}$$

(Here by abuse of notation, the arrows σ and ξ in the diagram are representatives of the points σ and ξ respectively.) If θ corresponds to a triple $(\lambda_R, (\overline{\lambda}_R, b_R), \text{id}_{\overline{\lambda}_R})$, then, the pull-backs $(\lambda_K, (\overline{\lambda}_K, b_K), \text{id}_{\overline{\lambda}_K})$ and $(\lambda_{\kappa}, (\overline{\lambda}_{\kappa}, b_{\kappa}), \text{id}_{\overline{\lambda}_{\kappa}})$ correspond to σ and ξ respectively. By extending R , we can assume that $\overline{\lambda}_R : \text{Spec } R \rightarrow \mathcal{X}$ lifts to $\nu_R : \text{Spec } R \rightarrow M$. Since p is étale, $\lambda_R : D_m \otimes R \rightarrow \mathcal{X}$ uniquely lifts to $\tilde{\nu} : D_m \otimes R \rightarrow M$ such that the diagram

$$\begin{array}{ccc} \text{Spec } R & \xrightarrow{\nu} & M \\ \downarrow & \nearrow \tilde{\nu} & \downarrow \\ D_m \otimes R & \xrightarrow{\lambda_R} & \mathcal{X} \end{array}$$

is commutative. Let $\overline{K}$ be the algebraic closure of K , let η be the composition $D_m \otimes \overline{K} \rightarrow D_m \otimes R \xrightarrow{\tilde{\nu}} M$ and let $b'_R : \text{Spec } R \rightarrow M \times_{\mathcal{X}} M$ be the lift of ν_R which corresponds to b_R . From ♣ and the assumption, there is a morphism $\delta : D_m \otimes \overline{K} \rightarrow M \times_{\mathcal{X}} M$ such that $\text{pr}_1 \circ \delta = \eta$ and $\text{pr}_2 \circ \delta = \eta \circ \zeta_l$, and the composition

$$\text{Spec } \overline{K} \hookrightarrow D_m \otimes \overline{K} \xrightarrow{\delta} M \times_{\mathcal{X}} M$$

equals the composition

$$\mathrm{Spec} \overline{K} \rightarrow \mathrm{Spec} R \xrightarrow{b'_R} M \times_{\mathcal{X}} M.$$

We can replace $\overline{K}$ with a finite extension K' of K . Moreover, replacing R with its normalization in K' , we can assume that $K' = K$ and that b'_R and δ induce the same morphism $\mathrm{Spec} K \rightarrow M \times_{\mathcal{X}} M$.

$$\begin{array}{ccccc} \mathrm{Spec} K & \longrightarrow & \mathrm{Spec} R & \xrightarrow{b'_R} & M \times_{\mathcal{X}} M \\ \downarrow & & \downarrow & \nearrow \delta & \downarrow \downarrow \\ D_m \otimes K & \longrightarrow & D_m \otimes R & \xrightarrow{\tilde{\nu}} & M \end{array}$$

Consider the unique morphism $\tau : D_m \otimes R \rightarrow M \times_{\mathcal{X}} M$ such that $\mathrm{pr}_1 \circ \tau = \tilde{\nu}$. Then the two morphism $\mathrm{pr}_2 \circ \tau$ and $\tau \circ \zeta_l$ is the same morphism because of the separatedness of M . Then the composition $D_m \otimes \kappa \rightarrow D_m \otimes R \xrightarrow{\tau} M \times_{\mathcal{X}} M$ satisfies the condition in $\clubsuit$. Hence $\xi \in \mathrm{Im}(\Psi)$. The proof is now complete. $\square$

Definition 3.6. We define the *stack of twisted n -jets of order l* on $\mathcal{X}$, denoted $\mathcal{L}_n^l \mathcal{X}$, to be the reduced closed substack of $L_m \mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})$ with support $\mathrm{Im}(\Psi)$. We define the *stack of twisted n -jets* on $\mathcal{X}$, denoted $\mathcal{L}_n \mathcal{X}$, to be the disjoint sum $\coprod_{l \geq 0} \mathcal{L}_n^l \mathcal{X}$. In particular, $\mathcal{L}_0 \mathcal{X}$ is the inertia stack $I(\mathcal{X})$.

If we set

$$l_0 := \max\{l \mid l = \mathrm{ord}(\alpha), \text{ for some } \xi \in \mathrm{ob} \mathcal{X} \text{ and for some } \alpha \in \mathrm{Aut}(\xi)\},$$

then for any $l > l_0$, $\mathcal{L}_n^l \mathcal{X} = \emptyset$. So the disjoint sum above is indeed a finite sum.

3.3. The formal neighborhood of $I(\mathcal{X})$ and its canonical automorphism. Let $\mathcal{X}$ be a smooth Deligne-Mumford stack, $x : \mathrm{Spec} k \rightarrow \mathcal{X}$ its closed point. Then the tangent space $T_x \mathcal{X}$ is defined to be $T_v M$ for an atlas $M \rightarrow \mathcal{X}$ and a lift $v : \mathrm{Spec} k \rightarrow M$ of x , uniquely determined up to unique isomorphism. Then $\mathrm{Aut}(x)$ naturally acts on $T_x \mathcal{X}$. We now globalize it.

Let $\mathcal{Y}$ be a connected component of the inertia stack $I(\mathcal{X})$, $F : \mathcal{Y} \rightarrow \mathcal{X}$ the forgetting map. Let $\mathcal{X}_0$ be the image of $\mathcal{Y}$ by F . The completion $\hat{\mathcal{O}}_{\mathcal{X}}$ of $\mathcal{O}_{\mathcal{X}}$ along $\mathcal{X}_0$ is considered to be an $\mathcal{O}_{\mathcal{X}_0}$ -algebra. Then we define a coherent sheaf $\mathcal{A}$ on $\mathcal{Y}$ to be the pull-back of $\hat{\mathcal{O}}_{\mathcal{X}}$ by $F : \mathcal{Y} \rightarrow \mathcal{X}_0$.

Definition 3.7. We define $\mathcal{N} := \mathrm{Spec} \mathcal{A}$ and call it the *formal neighborhood* of $\mathcal{Y}$.

If we set $\hat{\mathcal{X}} := \mathrm{Spec} \hat{\mathcal{O}}_{\mathcal{X}}$ where we consider $\hat{\mathcal{O}}_{\mathcal{X}}$ to be an $\mathcal{O}_{\mathcal{X}}$ -algebra, there is a natural morphism $\hat{\mathcal{X}} \rightarrow \mathcal{X}$, which is $\mathcal{X}_0$ -étale. Since $\mathcal{Y} \rightarrow \mathcal{X}_0$ is unramified and flat, it is étale. Hence the natural morphism $\mathcal{N} \rightarrow \hat{\mathcal{X}}$ is also étale and the composition $\mathcal{N} \rightarrow \hat{\mathcal{X}} \rightarrow \mathcal{X}$ is $\mathcal{Y}$ -étale.

Let U, V be varieties. Let $\xi : U \rightarrow \mathcal{X}$ be an étale morphism, $\nu : V \rightarrow U$ a morphism and α an automorphism of $\xi \circ \nu$. Suppose that $(\xi \circ \nu, \alpha) : V \rightarrow \mathcal{Y}$ is étale.

$$\begin{array}{ccc} V & \xrightarrow{\nu} & U \\ (\xi \circ \nu, \alpha) \downarrow & & \downarrow \xi \\ \mathcal{Y} & \longrightarrow & \mathcal{X} \end{array}$$

Then we obtain a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{\nu} & U \\ \nu \downarrow & \alpha \nearrow & \downarrow \xi \\ U & \xrightarrow{\xi} & \mathcal{X}, \end{array}$$

where ‘ $\Rightarrow$ ’ denotes a 2-morphism. Let $\tilde{\alpha} : V \rightarrow U \times_{\mathcal{X}} U$ be the corresponding morphism.

$$\begin{array}{ccc} & U \times_{\mathcal{X}} U & \\ \tilde{\alpha} \nearrow & \downarrow \text{pr}_1 & \downarrow \text{pr}_2 \\ V & \xrightarrow{\nu} & U. \end{array}$$

If $\hat{\mathcal{O}}_U$ is the completion of $\mathcal{O}_U$ along $\nu(V)$, then $\mathcal{A}_V = \nu^* \hat{\mathcal{O}}_U$. We have a canonical automorphism of $\mathcal{A}_V$

$$\mathcal{A}_V \xrightarrow{\text{pr}_2^*} \tilde{\alpha}^* \hat{\mathcal{O}}_{U \times_{\mathcal{X}} U} \xrightarrow{(\text{pr}_1^*)^{-1}} \mathcal{A}_V,$$

and hence a canonical automorphism of $\mathcal{A}$ and $\mathcal{N}$. Now this automorphism of $\mathcal{N}$ is considered to be a globalization of the action on $T_x \mathcal{X}$ mentioned above.

3.4. Shift number and orbifold cohomology. Suppose that $\mathcal{Y}$ is contained in $I^l(\mathcal{X})$ for an integer $l \geq 1$. Let (x, α) be a closed point of $\mathcal{Y}$ where x is a closed point of $\mathcal{X}$ and $\alpha \in \text{Aut}(x)$. Then α acts on the tangent space $T_x \mathcal{X}$. For a suitable basis, this automorphism is given by a diagonal matrix

$$\text{diag}(\zeta_l^{a_1}, \dots, \zeta_l^{a_d})$$

with $1 \leq a_j \leq l$ and $d = \dim \mathcal{X}$.

Definition 3.8. We define the *shift number* of $\mathcal{Y}$ by

$$s(\mathcal{Y}) := \dim \mathcal{X} - \frac{1}{l} \sum_{j=1}^d a_j = \frac{1}{l} \sum_{j=1}^d (l - a_j).$$

This is determined by the rank of the eigenbundles of $\mathcal{N}$ for the canonical action. Hence it depends only on $\mathcal{Y}$.

Suppose that the coarse moduli space $X = \overline{\mathcal{X}}$ is a variety with Gorenstein quotient singularities and $\mathcal{X}$ has no reflections. Then the matrix $\text{diag}(\zeta_l^{a_1}, \dots, \zeta_l^{a_d})$ is in $\text{SL}_d(\mathbb{C})$ (see [Wat74]). Hence $s(\mathcal{Y})$ is an integer.

Now, let us define the orbifold cohomology.

Definition 3.9. Assume X is complete. Then we define the i -th orbifold cohomology group along with Hodge structure as follows:

$$H_{\text{orb}}^i(X, \mathbb{Q}) := \bigoplus_{\mathcal{Y}} H^{i-2s(\mathcal{Y})}(\overline{\mathcal{Y}}, \mathbb{Q}) \otimes \mathbb{Q}(-s(\mathcal{Y})),$$

where $\mathcal{Y}$ runs over the connected components of $I(\mathcal{X})$ and $\mathbb{Q}(-s(\mathcal{Y}))$ is a Tate twist $\mathbb{Q}(-1)^{\otimes s(\mathcal{Y})}$.

Since the natural morphism $\overline{\mathcal{Y}} \rightarrow X$ is quasi-finite, $\overline{\mathcal{Y}}$ is a scheme (see [LMB00, Th. A.2]). Because of this and Corollary 4.23, $\overline{\mathcal{Y}}$ is a complete variety with quotient singularities. Therefore the rational cohomology groups of $\mathcal{Y}$ have *pure* Hodge structures. For the projective case, see [Dan78, Cor. 14.4]. For the general case, it follows from the following two facts: one is that the intersection cohomology of every complete variety has pure Hodge structures [Sai90], the other is that the rational cohomology of a variety with quotient singularities equals the intersection cohomology.

3.5. The motivic measure on twisted ∞ -jets. Let $\mathcal{X}$ be a smooth Deligne-Mumford stack of pure dimension d . By abuse of notation, we also denote by $\overline{\mathcal{L}_\infty \mathcal{X}}$ the set of points $|\mathcal{L}_\infty \mathcal{X}| = |\overline{\mathcal{L}_\infty \mathcal{X}}|$. We denote by π_n the natural morphism $\overline{\mathcal{L}_\infty \mathcal{X}} \rightarrow \overline{\mathcal{L}_n \mathcal{X}}$.

Definition 3.10. A subset A of $\overline{\mathcal{L}_\infty \mathcal{X}}$ is *stable at level n* if we have:

- (1) $\pi_n(A)$ is a constructible subset in $\overline{\mathcal{L}_n \mathcal{X}}$,
- (2) $A = \pi_n^{-1}\pi_n(A)$,

A subset $A \subset \overline{\mathcal{L}_\infty \mathcal{X}}$ is *stable* if it is stable at level n for some $n \in \mathbb{Z}_{\geq 0}$.

We define the notion of the *measurable subset* similarly. Then we define *motivic measure* $\mu_{\mathcal{X}}$ on $\overline{\mathcal{L}_\infty \mathcal{X}}$ by

$$\mu_{\mathcal{X}}(A) := \mathbb{L}^{-nd} \chi_h(\pi_n(A)) \in \hat{K}_0(\text{HS}), \quad n \gg 0,$$

where $\mathbb{L} = \{\mathbb{Q}(-1)\} = \chi_h(\mathbb{A}^1)$. It is well-defined by the following:

Proposition 3.11. Let $n \in \mathbb{Z}_{\geq 0}$, let $B \subset \overline{\mathcal{L}_n \mathcal{X}}$ be a constructible subset and let C be the inverse image of B by the natural morphism $\overline{\mathcal{L}_{n+1} \mathcal{X}} \rightarrow \overline{\mathcal{L}_n \mathcal{X}}$. Then we have the equality $\chi_h(C) = \mathbb{L}^d \chi_h(B)$.

Proof. Let $\mathcal{Y}$ be a connected component of $I^l(\mathcal{X}) (= \mathcal{L}_0^l \mathcal{X})$ and put $(\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} := \mathcal{L}_n \mathcal{X} \times_{I(\mathcal{X})} \mathcal{Y}$. Let $\gamma = (\tilde{\gamma}, (\bar{\gamma}, b), \text{id}_{\bar{\gamma}}) \in (\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \subset \mathcal{L}_m \mathcal{X} \times_{\mathcal{X}} \mathcal{Y}$ be an Ω -point. Then we have the following commutative diagram of solid arrows:

$$\begin{array}{ccc} D_0 \otimes \Omega & \xrightarrow{(\bar{\gamma}, b)} & \mathcal{Y} \longrightarrow \mathcal{N} \\ \downarrow & \nearrow \sigma & \downarrow \\ D_m \otimes \Omega & \xrightarrow{\tilde{\gamma}} & \mathcal{X}. \end{array}$$

Since $\mathcal{N} \rightarrow \mathcal{X}$ is $\mathcal{Y}$ -étale, there is a unique broken arrow σ fitting into the diagram. Sending γ to σ determines a closed immersion

$$\iota : (\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \hookrightarrow L_m \mathcal{N}.$$

Let $\mathbf{g}$ be the canonical automorphism of $\mathcal{N}$. In view of the definition of $\mathcal{N}$ and $\clubsuit$ in the proof of Lemma 3.5, we see that for $\sigma \in L_m \mathcal{N}$, $\sigma \in \text{Im}(\iota)$ iff $\sigma \circ \zeta_l = \mathbf{g} \circ \sigma$. From Lemma 3.12, there exists an atlas $h : V \rightarrow \mathcal{Y}$ such that $\mathcal{N}_V := \mathcal{N} \times_{\mathcal{Y}} V \cong V \otimes \mathbb{C}[[v_1, \dots, v_c]]$, ($c = \dim \mathcal{X} - \dim \mathcal{Y}$), and the pull-back of $\mathbf{g}$ is given by $\text{diag}(\zeta_l^{a_1}, \dots, \zeta_l^{a_c})$, $1 \leq a_i \leq l$. So for an m -jet δ on $\mathcal{N}_V$, $h \circ \delta$ is in $\text{Im}(\iota)$ iff the image of δ by $L_m(\mathcal{N}_V) \rightarrow L_m V$ is ζ_l -invariant and $\delta^*(v_i)$ is of the following form:

$$r_0 t^{a_i} + r_1 t^{a_i+l} + r_2 t^{a_i+2l} + \dots.$$

Therefore we have that $(\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \times_{\mathcal{Y}} V \cong L_n V \times \mathbb{A}^{nc}$ and the projection $(\mathcal{L}_{n+1} \mathcal{X})_{\mathcal{Y}} \times_{\mathcal{Y}} V \rightarrow (\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \times_{\mathcal{Y}} V$ is a Zariski locally trivial $\mathbb{A}^d$ -bundle.

We may assume that a finite group, say G , acts on each connected component V' of V , $V' \rightarrow \overline{\mathcal{Y}}$ is G -invariant and the induced morphism $V'/G \rightarrow \overline{\mathcal{Y}}$ is étale (see Lemma 4.26). Then we have that $(\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \times_{\overline{\mathcal{Y}}} (V'/G) \cong ((L_n V')/G) \times \mathbb{A}^{nc}$. Therefore, from Lemma 3.14, to prove the proposition, it suffices to show that there is a stratification of $(L_n V')/G$ such that the natural morphism $(L_{n+1} V')/G \rightarrow (L_n V')/G$ is, over each stratum, an analytically locally trivial fibration of the quotient of an affine space by a linear finite group action.

Let H be a subgroup of G and $W \subset V'$ a connected component of the locus of the points with stabilizer H . Let $w \in W$ be a close point. As is well known, there is a representation $\rho : H \subset \text{GL}_{\dim \mathcal{Y}}(\mathbb{C})$ which describes the H -action on an analytic neighborhood of w . Let $(L_n V)_w \subset L_n V$ be the subset of the jets which maps the only point of D_n to w . Then the induced H -action on $(L_n V)_w \cong \mathbb{A}^{m \dim \mathcal{Y}}$ is given by $\rho^{\oplus m}$. Therefore $(L_{n+1} V')_W/H \rightarrow (L_n V')_W/H$ is an analytically locally trivial fibration of $\mathbb{A}^{\dim \mathcal{Y}}/H$. Let $G' \subset G$ be the subgroup of the elements keeping W stable. Then H is a normal subgroup of G' . It is easy to see that the image of $(L_n V')_W/H$ in $(L_n V')/G$ is naturally isomorphic to $((L_n V')_W/H)/(G'/H)$. Since G'/H freely acts on $(L_n V')_W/H$, the assertion follows. $\square$

Lemma 3.12. *Let G be a finite group, V a smooth G -variety and W a smooth closed subvariety consisting of G -invariant points.*

- (1) *Assume that V and W are affine, say $V = \text{Spec } R$ and $W = \text{Spec } R/\mathfrak{p}$. Moreover assume that $\mathfrak{p}$ is generated by $c = \text{codim}(W, V)$ elements. Then the completion of V along W is isomorphic as G -schemes to $\text{Spec}(R/\mathfrak{p})[[x_1, \dots, x_c]]$, $G \subset \text{GL}_c(R/\mathfrak{p})$.*
- (2) *Assume that G is a finite cyclic group. Then there is an affine open covering $\cup V_i$ of V such that for every i , if we write $V_i = \text{Spec } R$ and $W \cap V_i = \text{Spec } R/\mathfrak{p}$, the completion of V_i along $W \cap V_i$ is isomorphic as G -schemes to $\text{Spec}(R/\mathfrak{p})[[x_1, \dots, x_c]]$, $G \subset \text{GL}_c(\mathbb{C})$.*

Proof. (1). We denote by $\hat{R}$ the completion of R with respect to an ideal $\mathfrak{p}$ and by $\widehat{R_{\mathfrak{p}}}$ the completion of the local ring $R_{\mathfrak{p}}$ with respect to the maximal ideal. Let K be the quotient field of $R/\mathfrak{p}$ and f_i generators of $\mathfrak{p}$. It is well known that there is an isomorphism $\widehat{R_{\mathfrak{p}}} \rightarrow K[[x_1, \dots, x_c]]$ sending f_i to x_i . We have a natural injection $\hat{R} \rightarrow \widehat{R_{\mathfrak{p}}} \cong K[[x_1, \dots, x_c]]$. Clearly the image contains the subring $(R/\mathfrak{p})[[x_1, \dots, x_c]]$. Consider the injection $\iota : (R/\mathfrak{p})[[x_1, \dots, x_c]] \rightarrow \hat{R}$. Since the induced map $R/\mathfrak{p} \rightarrow \hat{R}/\hat{\mathfrak{p}}$ is the identity and the images f_i of x_i generate $\hat{\mathfrak{p}}$, ι is a surjection and hence an isomorphism (see [Eis95, Th. 7.16.]).

Consider the induced G -action on $\hat{R} = (R/\mathfrak{p})[[x_1, \dots, x_c]]$. For $g \in G$, write $g(x_i) = \sum a_{ij}x_j + (\text{higher terms})$, $a_{ij} \in R/\mathfrak{p}$. Denote by $\bar{g}$ the endomorphism of $\hat{R}$ associated to the invertible matrix (a_{ij}) . Working in characteristic zero, we can isomorphically replace x_i with $x'_i = \sum_{g \in G} \bar{g}^{-1}g(x_i)$. We find that with respect to the new coordinates, the G -action is linear.

(2). In the last situation, $\text{Spec } \hat{R}$ is naturally isomorphic to the completion of the normal bundle $N_{W/V}$ along the zero section. Here let us assume G is a cyclic group with generator g . Then $N_{W/V}$ decomposes to eigenbundles. On each eigenbundle, the g -action is uniquely represented by a scalar matrix aI where I is the identity matrix and $a \in \mathbb{C}$. Therefore shrinking V to an open subset where the eigenbundles are free, we conclude that the G -actions on $N_{W/V}$ and $\text{Spec } \hat{R}$ are realizable in $\mathbb{C}$. $\square$

Remark 3.13. The author guesses that even in the case of a general finite group, the action on $N_{W/V}$ is *étale* locally realizable in $\mathbb{C}$. From facts on splitting fields of finite groups (see [CR88]), this is true at least over the generic point of W .

Lemma 3.14. *Let T and S be varieties and $f : T \rightarrow S$ an analytically locally trivial fibration of $\mathbb{A}^d/G$ for a finite group $G \subset \text{GL}_d(\mathbb{C})$. Then $\chi_h(T) = \chi_h(S)\mathbb{L}^d$.*

Proof. Since the fiber is a quotient of an affine space, the higher direct images of $\mathbb{Q}_T$ vanishes;

$$R^i f_* \mathbb{Q}_T \cong \begin{cases} \mathbb{Q}_S & (i = 0) \\ 0 & (i > 0). \end{cases}$$

Hence the spectral sequence is degenerate and it follows that $H^i(T, \mathbb{Q}) \cong H^i(S, \mathbb{Q})$ for every i .

Taking a stratification of S , we may assume that S is smooth. Since S and T have at most quotient singularities (in the analytic sense), by Poincaré duality, we conclude that $H_c^{\dim T - i}(T, \mathbb{Q}) \cong H_c^{\dim S - i}(S, \mathbb{Q})$.

Regarding the sheaves as mixed Hodge modules, studied by Saito [Sai90] (see also [Sai89]), we can regard the isomorphisms above of cohomology groups as ones of mixed Hodge structures. This implies the assertion. $\square$

3.6. Main theorem. Let X be a variety with Gorenstein quotient singularities. Then X has canonical singularities. Let $\mathcal{X}$ be a smooth Deligne-Mumford stack without reflections such that X is the coarse moduli space of $\mathcal{X}$. We denote by λ the canonical morphism $\mathcal{X} \rightarrow X$. If $\gamma : \mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$ is a twisted n -jet on $\mathcal{X}$ of order l , then it induces a morphism $\gamma' : D_n \otimes \Omega \rightarrow X$ of the coarse moduli spaces. We define the map $\lambda_{(n)} : \overline{\mathcal{L}_n \mathcal{X}} \rightarrow L_n X$ by $\gamma \mapsto \gamma'$. The following is our main result.

Theorem 3.15. *Let $B \subset \overline{\mathcal{L}_\infty \mathcal{X}}$ be a measurable subset and put $A := \lambda_{(\infty)}(B)$. Then we have the following equation in $\hat{K}_0(\text{HS})$:*

$$\chi_h \mu_X^{\text{Gor}}(A) = \sum_{\mathcal{Y} \subset I(\mathcal{X})} \mathbb{L}^{s(\mathcal{Y})} \mu_{\mathcal{X}}(\pi_0^{-1}(\overline{\mathcal{Y}}) \cap B),$$

where $\mathcal{Y}$ runs over the connected components of $I(\mathcal{X})$.

The proof is postponed until the end of the section.

Corollary 3.16. *Let X and X' be complete varieties with Gorenstein quotient singularities. Suppose that there are proper birational morphisms $Z \rightarrow X$ and $Z \rightarrow X'$ such that $K_{Z/X} = K_{Z/X'}$. Then the orbifold cohomology groups of X and X' have the same Hodge structure.*

Proof. By Theorem 3.15 and Proposition 3.11, we have

$$\begin{aligned} \chi_h \mu_X^{\text{Gor}}(L_\infty X) &= \sum_{\mathcal{Y} \subset I(\mathcal{X})} \mathbb{L}^{s(\mathcal{Y})} \chi_h(\overline{\mathcal{Y}}) \\ &= \sum_i (-1)^i \{H_{\text{orb}}^i(X, \mathbb{Q})\}. \end{aligned}$$

From Lemma 2.16, we have $\sum_i (-1)^i \{H_{\text{orb}}^i(X, \mathbb{Q})\} = \sum_i (-1)^i \{H_{\text{orb}}^i(X', \mathbb{Q})\}$. Since $H_{\text{orb}}^i(X, \mathbb{Q})$ and $H_{\text{orb}}^i(X', \mathbb{Q})$ have a pure Hodge structure of weight i , $\{H_{\text{orb}}^i(X, \mathbb{Q})\} = \{H_{\text{orb}}^i(X', \mathbb{Q})\}$ for every i . $\square$

Lemma 3.17. *Let X_{sing} denote the singular locus of X with reduced subscheme structure.*

- (1) *The subset $\lambda_{(\infty)}^{-1}(L_\infty(X_{\text{sing}}))$ of $\overline{\mathcal{L}_\infty \mathcal{X}}$ is of measure zero.*
- (2) *The map $\lambda_{(\infty)}$ is bijective over $L_\infty X \setminus L_\infty(X_{\text{sing}})$.*

Proof. (1). It suffices to show that for every n , $\pi_n(\overline{\mathcal{L}_\infty \mathcal{X}} \setminus \lambda_{(\infty)}^{-1}(L_\infty(X_{\text{sing}}))) = \overline{\mathcal{L}_n \mathcal{X}}$. But this is clear by the local description of $\overline{\mathcal{L}_n \mathcal{X}}$ in the proof of Proposition 3.11.

(2). Surjectivity. Let $\eta : \text{Spec } \Omega[[t]] \rightarrow X$ be an Ω -point of $L_\infty X \setminus L_\infty(X_{\text{sing}})$ with an algebraically closed field Ω . We define $\mathcal{D}$ to be the normalization of the fiber product $\mathcal{X} \times_X \text{Spec } \Omega[[t]]$, (see Definition 4.13). Then the Deligne-Mumford stack $\mathcal{D}$ contains the scheme $\text{Spec } \Omega((t))$ as open substack. Therefore the coarse moduli space $\overline{\mathcal{D}}$ of $\mathcal{D}$ contains $\text{Spec } \Omega((t))$ as open subscheme. The scheme $\overline{\mathcal{D}}$ must be the spectrum of a local ring $S \subset \Omega((t))$. From the universality of coarse moduli space, there is a natural

morphism $\overline{\mathcal{D}} \rightarrow \operatorname{Spec} \Omega[[t]]$. So we have $\Omega[[t]] \subset S \subset \Omega((t))$. Because $\Omega[[t]]$ and $\Omega((t))$ are the only intermediate rings between $\Omega[[t]]$ and $\Omega((t))$, the ring S must be $\Omega[[t]]$. Suppose that $E = \operatorname{Spec} \tilde{S}$ is an atlas of $\mathcal{D}$ and $\tilde{S}$ is a regular local ring. Since $S = \Omega[[t]]$ is henselian, the natural morphism $E \rightarrow \overline{\mathcal{D}}$ is finite ([EGA, IV. Th. 18.5.11]). Hence $\tilde{S}$ is complete, (see [Eis95, Cor. 7.6]). So $\tilde{S} \cong \Omega[[t]]$. Consider the groupoid space $E \times_{\mathcal{D}} E \rightrightarrows E$. The scheme $E \times_{\mathcal{D}} E$ must be the disjoint sum of spectra of complete regular local rings. Since the first projection $\operatorname{pr}_1 : E \times_{\mathcal{D}} E \rightarrow E$ is étale, there is an isomorphism

$$E \coprod \cdots \coprod E \cong E \times_{\mathcal{D}} E$$

such that the composition

$$E \coprod \cdots \coprod E \cong E \times_{\mathcal{D}} E \xrightarrow{\operatorname{pr}_1} E$$

is isomorphic on each component. If l denote the number of the components in $E \times_{\mathcal{D}} E$, then the second projection $\operatorname{pr}_2 : E \times_{\mathcal{D}} E \rightarrow E$ determines the action of some group G on E with $|G| = l$. Since this action is effective, the group G is isomorphic to μ_l for some l . For a suitable isomorphism $\mu_l \cong G$, the action is given by $t \mapsto \zeta_l t$. Hence the stack $\mathcal{D}$ is isomorphic to $\mathcal{D}_{\infty}^l \otimes \Omega$ and the morphism $\mathcal{D}_{\infty}^l \otimes \Omega \cong \mathcal{D} \rightarrow \mathcal{X}$ is a twisted ∞ -jet on $\mathcal{X}$. The image of this twisted ∞ -jet by $\lambda_{(\infty)}$ is η .

Injectivity. Let $\gamma_1, \gamma_2 : \mathcal{D}_{\infty}^l \otimes \Omega \rightarrow \mathcal{X}$ be two twisted ∞ -jets on $\mathcal{X}$ of order l . We suppose that $\eta := \lambda_{(\infty)}(\gamma_1) = \lambda_{(\infty)}(\gamma_2)$ and $\eta \in L_{\infty} X \setminus L_{\infty}(X_{\text{sing}})$. Construct $\mathcal{D}$ from η as above. Then for each $i \in \{1, 2\}$, there is a unique morphism $h_i : \mathcal{D}_{\infty}^l \otimes \Omega \rightarrow \mathcal{D}$ such that the following diagram is commutative:

$$\begin{array}{ccccc} \operatorname{Spec} \Omega((t)) & \hookrightarrow & \mathcal{D}_{\infty}^l \otimes \Omega & & \\ \parallel & & \downarrow h_i & \searrow \gamma_i & \\ \operatorname{Spec} \Omega((t)) & \hookrightarrow & \mathcal{D} & \longrightarrow & \mathcal{X} \\ \parallel & & \downarrow & & \downarrow \lambda \\ \operatorname{Spec} \Omega((t)) & \hookrightarrow & D_{\infty} \otimes \Omega & \xrightarrow{\eta} & X. \end{array}$$

Let E be an atlas of $\mathcal{D}$ as above. Then the natural morphism $E \times_{\mathcal{D}, h_i} (\mathcal{D}_{\infty}^l \otimes \Omega) \rightarrow E$ is a birational morphism of smooth 1-dimensional schemes. Therefore it is an isomorphism and so is h_i ([LMB00, Prop. 3.8.1]). Then we can easily see that γ_1 and γ_2 have the same image in $|L_{\infty} \mathcal{X} \times_{\mathcal{X}} I^l(\mathcal{X})|$. $\square$

To prove Theorem 3.15, we now need to generalize the transformation rule. Let $\mathcal{V}$ be a Deligne-Mumford stack over D_{∞} of pure relative dimension d . For each $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$, we define $\mathcal{V}_n$ to be the moduli stack of the D_{∞} -morphisms $D_n \rightarrow \mathcal{V}$. Then for $m \geq n$, there is a natural projection $\mathcal{V}_m \rightarrow \mathcal{V}_n$. So we can define the *motivic measure* $\mu_{\mathcal{V}}$ over $\overline{\mathcal{V}_{\infty}}$ which takes values in $\hat{K}_0(\text{HS})$, in a similar fashion as before. (We should replace condition (3) in

Definition 2.7 with the condition that $\chi_h(\pi_{m+1}A) = \mathbb{L}^d \chi_h(\pi_m A)$. It makes sense because of Lemma 3.14.)

Let $\mathcal{W}$ be another Deligne-Mumford stack over D_∞ of pure relative dimension d and let $h : \mathcal{W} \rightarrow \mathcal{V}$ be a D_∞ -morphism. We put $\Omega'_{\mathcal{W}/\mathcal{V}} := \text{Im}(\Omega_{\mathcal{W}/D_\infty} \setminus (\text{tors}) \rightarrow \Omega_{\mathcal{W}/\mathcal{V}})$, where $(\text{tors}) \subset \Omega_{\mathcal{W}/D_\infty}$ is the torsion. Then we define the *jacobian ideal sheaf* $\mathcal{J}_h$ of h to be the 0-th Fitting ideal of $\Omega'_{\mathcal{W}/\mathcal{V}}$.

Theorem 3.18. *Let $A \subset \overline{\mathcal{W}_\infty}$ be a measurable set. Suppose that $h_\infty|_A : A \rightarrow \overline{\mathcal{V}_\infty}$ is injective. Let ν be a measurable function on $h_\infty(A)$. Then*

$$\int_{h_\infty(A)} \mathbb{L}^\nu d\mu_{\mathcal{V}} = \int_A \mathbb{L}^{\nu \circ h_\infty - \text{ord } \mathcal{J}_h} d\mu_{\mathcal{W}}.$$

Proof. It is a direct consequence of the following lemma. $\square$

We denote by $\mathcal{J}(\mathcal{V}/D_\infty)$ (resp. $\mathcal{J}(\mathcal{W}/D_\infty)$) the d -th Fitting ideal sheaf of $\Omega_{\mathcal{V}/D_\infty}$ (resp. $\Omega_{\mathcal{W}/D_\infty}$).

Lemma 3.19. *Let $A \subset \overline{\mathcal{W}_\infty}$ be a stable subset of level l . Assume that $h_\infty|_A$ is injective, that $\text{ord } \mathcal{J}_h$ is constant equal to $e < \infty$ and that $\text{ord } \mathcal{J}(\mathcal{V}/D_\infty)$ and $\text{ord } \mathcal{J}(\mathcal{W}/D_\infty)$ are bounded from above on $h_\infty(A)$ and A respectively. Then for $n \gg 0$, $h_n : \pi_n A \rightarrow h_n \pi_n A$ is a piecewise trivial $\mathbb{A}^e$ -bundle.*

Proof. Looijenga's proof [Loo02, Lem. 9.2] works also in this setting.

Take a non-twisted ∞ -jet $\gamma : \text{Spec } \Omega[[t]] \rightarrow \mathcal{W}$ in A , and put $\mathfrak{m} := (t) \subset \Omega[[t]]$. Let q be the image of the closed point by γ . Take another $\theta \in A$ such that $\pi_{n-e}(\gamma) = \pi_{n-e}(\theta)$. Then the morphism

$$\theta^* - \gamma^* : \mathcal{O}_{\mathcal{W},q} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{2(n-e+1)}$$

is a $\mathbb{C}[[t]]$ -derivation. So it defines an $\Omega[[t]]$ -module homomorphism

$$\partial\theta : \gamma^* \Omega_{\mathcal{W}/D_\infty} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{2(n-e+1)}.$$

The length of the torsion of $\gamma^* \Omega_{\mathcal{W}/D_\infty}$ equals $(\text{ord } \mathcal{J}(\mathcal{W}/D_\infty))(\gamma)$, hence it is bounded. So, since $n \gg 0$, the composition map

$$\overline{\partial\theta} : \gamma^* \Omega_{\mathcal{W}/D_\infty} \xrightarrow{\partial\theta} \mathfrak{m}^{n-e+1}/\mathfrak{m}^{2(n-e+1)} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1}$$

annihilates the torsion. Conversely, every $\Omega[[t]]$ -module homomorphism $\gamma^* \Omega_{\mathcal{W}/D_\infty} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1}$ which annihilates the torsion is $\overline{\partial\theta}$ for some θ .

After some works, we can see that if $\theta \in A$ is such that $h_n \pi_n(\gamma) = h_n \pi_n(\theta)$, then $\pi_{n-e}(\gamma) = \pi_{n-e}(\theta)$, (see [Loo02, Lem. 9.2]). So $\overline{\partial\theta}$ is defined. It is easy to see that $\pi_n(\theta_1) = \pi_n(\theta_2)$ iff $\overline{\partial\theta_1} = \overline{\partial\theta_2}$, and that $h_n \pi_n(\theta_1) = h_n \pi_n(\theta_2)$ iff $\overline{\partial\theta_1}$ and $\overline{\partial\theta_2}$ have the same image in $\text{Hom}_{\Omega[[t]]}((h\gamma)^* \Omega_{\mathcal{V}/D_\infty}, \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1})$.

Hence $h_n^{-1}h_n\pi_n(\gamma)$ is isomorphic to an affine space,

$$\begin{aligned} \mathrm{Hom}_{\Omega[[t]]}(\gamma^*\Omega'_{\mathcal{W}/\mathcal{V}}, \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1}) \\ \cong \mathrm{Ker}(\mathrm{Hom}_{\Omega[[t]]}(\gamma^*\Omega_{\mathcal{W}/D_\infty} \setminus (tors), \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1}) \\ \rightarrow \mathrm{Hom}_{\Omega[[t]]}((h\gamma)^*\Omega_{\mathcal{V}/D_\infty}, \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1})). \end{aligned}$$

The length of $\gamma^*\Omega'_{\mathcal{W}/\mathcal{V}}$ equals $e = \mathrm{ord} \mathcal{J}_h(\gamma)$. So $h_n^{-1}h_n\pi_n(\gamma)$ is isomorphic to an affine space of dimension e .

The rest is easy. $\square$

Proof of Theorem 3.15 . Let $\mathcal{Y}$ be a connected component of $I^l(\mathcal{X})$ and $\mathcal{N}$ its formal neighborhood. We may assume that B is contained in $\pi_0^{-1}(\overline{\mathcal{Y}})$. Let $\tilde{\mathcal{N}}$ be the quotient of $\mathcal{N}$ by the canonical automorphism $\mathfrak{g}$, that is, $\mathrm{Spec} \mathcal{A}^{\mathfrak{g}}$ where $\mathcal{A}^{\mathfrak{g}} \subset \mathcal{A}$ is the subsheaf of the $\mathfrak{g}$ -invariant sections. Then the natural morphism $\mathcal{N} \rightarrow X$ factors as

$$\mathcal{N} \rightarrow \tilde{\mathcal{N}} \xrightarrow{f} X.$$

In the proof of Proposition 3.11, we saw that for each n, m with $m = nl$, there is a closed immersion $\iota : (\mathcal{L}_n \mathcal{X})_{\mathcal{Y}} \hookrightarrow L_m \mathcal{N}$. Let $[l] : D_m \rightarrow D_n$ be the morphism associated to the ring homomorphism defined by $t \mapsto t^l$. If $\gamma : \mathcal{D}_n^l \otimes \Omega \rightarrow \mathcal{X}$ is a twisted n -jet in $\pi_n(B)$, then $\iota(\gamma)$ fits into the diagram

$$(3.1) \quad \begin{array}{ccc} D_m \otimes \Omega & \xrightarrow{\iota(\gamma)} & \mathcal{N} \\ [l] \downarrow & & \downarrow \\ D_n \otimes \Omega & \xrightarrow{\sigma} & \tilde{\mathcal{N}}. \end{array}$$

Then $\lambda_{(n)}(\gamma) = f_n(\sigma)$. We define a subset $\tilde{B} \subset \overline{L_\infty \tilde{\mathcal{N}}}$ to be the image of B by the map $\gamma \mapsto \sigma$. Then $A = f_\infty(\tilde{B})$ and $f_\infty|_{\tilde{B}}$ is bijective outside of subsets of measure zero. Let $\mathcal{J}_X$ (resp. $\mathcal{J}_{\tilde{\mathcal{N}}}$) be the ideal sheaf on X defined by

$$\begin{aligned} \mathcal{J}_X \omega_X &= \mathrm{Im}(\Omega_X^d \rightarrow \omega_X) \\ (\text{resp. } \mathcal{J}_{\tilde{\mathcal{N}}} \omega_{\tilde{\mathcal{N}}} &= \mathrm{Im}(\Omega_{\tilde{\mathcal{N}}}^d \rightarrow \omega_{\tilde{\mathcal{N}}}), \end{aligned}$$

and define μ_X^{Gor} and $\mu_{\tilde{\mathcal{N}}}^{\mathrm{Gor}}$ to be $\mathbb{L}^{\mathrm{ord} \mathcal{J}_X} \mu_X$ and $\mathbb{L}^{\mathrm{ord} \mathcal{J}_{\tilde{\mathcal{N}}}} \mu_{\tilde{\mathcal{N}}}$ respectively. Since the morphism f has no ramification divisor, by a similar argument as the proof of Lemma 2.16, we see $f^{-1}\mathcal{J}_X = \mathcal{J}_f \cdot \mathcal{J}_{\tilde{\mathcal{N}}}$, where $\mathcal{J}_f$ is the jacobian ideal sheaf. So, by Theorem 3.18, we obtain

$$\mu_{\tilde{\mathcal{N}}}^{\mathrm{Gor}}(\tilde{B}) = \mu_X^{\mathrm{Gor}}(A).$$

We have thus reduced the problem to the case of a cyclic quotient; it suffices to show the following lemma. $\square$

Lemma 3.20. *Let the notation as above. We have $\mathbb{L}^{s(\mathcal{Y})} \mu_{\mathcal{X}}(B) = \chi_h \mu_{\tilde{\mathcal{N}}}^{\mathrm{Gor}}(\tilde{B})$.*

Proof. The proof is essentially by a trick used in [DL02]. We first consider an easy case where $\mathcal{X}$ is a quotient stack $[\mathbb{A}_R^c/G]$ of an affine space over a ring R whose spectrum is a smooth variety of dimension $d - c$, and $G \subset \mathrm{SL}_c(\mathbb{C})$ a finite cyclic group of order l generated by $g = \mathrm{diag}(\zeta_l^{a_1}, \dots, \zeta_l^{a_c})$, $1 \leq a_i < l$. Suppose that $\mathcal{Y}$ is the component associated to g . Then $\mathcal{N} = \hat{\mathbb{A}}_R^c (= \mathrm{Spec} R[[x_1, \dots, x_c]])$, its canonical automorphism is $g = \mathrm{diag}(\zeta_l^{a_1}, \dots, \zeta_l^{a_c})$, and $\tilde{\mathcal{N}} = \hat{\mathbb{A}}_R^c/G$. Since the natural morphisms $\hat{\mathbb{A}}_R^c \rightarrow \mathbb{A}_R^c$ and $\hat{\mathbb{A}}_R^c/G \rightarrow \mathbb{A}_R^c/G$ are $(\mathrm{Spec} R)$ -étale, and since we consider only jets which send the only closed point into $\mathrm{Spec} R$, it makes no matter to replace $\hat{\mathbb{A}}_R^c$, $\hat{\mathbb{A}}_R^c/G$ with $\mathbb{A}_R^c$, $\mathbb{A}_R^c/G$.

Consider three R -algebra homomorphisms; (i) $u^* : R[[t]][\underline{x}] \rightarrow R[[t]][\underline{x}]$, $x_i \mapsto t^{a_i} x_i$, (ii) $\alpha^* : R[[t]][\underline{x}] \rightarrow R[[t]][\underline{x}]$, $x_i \mapsto x_i$, $t \mapsto t^l$, (iii) $\beta^* : R[[t]][\underline{x}]^G \rightarrow R[[t]][\underline{x}]$, the composition of α^* and the inclusion $R[[t]][\underline{x}]^G \hookrightarrow R[[t]][\underline{x}]$. Since $R[\underline{x}]^G$ is generated by the monomials $x_1^{m_1} \dots x_c^{m_c}$ with $\sum a_i m_i \equiv 0 \pmod{l}$, there is a $R[[t]]$ -homomorphism $v^* : R[[t]][\underline{x}]^G \rightarrow R[[t]][\underline{x}]$ with $u^* \circ \beta^* = \alpha^* \circ v^*$;

$$\begin{array}{ccc} R[[t]][\underline{x}] & \xrightarrow{u^*} & R[[t]][\underline{x}] \\ \beta^* \uparrow & & \uparrow \alpha^* \\ R[[t]][\underline{x}]^G & \xrightarrow{v^*} & R[[t]][\underline{x}]. \end{array}$$

Here the horizontal arrows are $R[[t]]$ -algebra homomorphisms and the vertical ones send $t \mapsto t^l$. Write the diagram of the associated schemes as follows;

$$\begin{array}{ccc} \mathcal{N}[[t]] & \xleftarrow{u} & E_2[[t]] \\ \beta \downarrow & & \downarrow \alpha \\ \tilde{\mathcal{N}}[[t]] & \xleftarrow{v} & E_1[[t]]. \end{array}$$

Here E_i are copies of $\mathbb{A}_R^c$.

Let B_0 be the image of B by $\iota : (\mathcal{L}_\infty \mathcal{X})_{\mathcal{Y}} \hookrightarrow L_\infty \mathcal{N}$. Then for $\gamma' \in B_0$, $(\gamma')^*(x_i)$ is of the form,

$$(3.2) \quad (\gamma')^*(x_i) = r_0 t^{a_i} + r_1 t^{a_i+l} + r_2 t^{a_i+2l} + \dots.$$

If we put $\eta := u_\infty^{-1}(\gamma')$, then we have

$$\eta^*(x_i) = r_0 + r_1 t^l + r_2 t^{2l} + \dots.$$

Therefore if we define $\delta \in L_\infty E_1$ by

$$(3.3) \quad \delta^*(x_i) = r_0 + r_1 t + r_2 t^2 + \dots,$$

then we have the following commutative diagram,

$$\begin{array}{ccc} E_2[[t]] & \xleftarrow{\eta} & D_\infty \otimes \Omega \\ \alpha \downarrow & & \downarrow [l] \\ E_1[[t]] & \xleftarrow[\delta]{} & D_\infty \otimes \Omega. \end{array}$$

Here $[l]$ is the morphism defined by $t \mapsto t^l$. Let $B_1 \subset L_\infty(E_1)$ be the image of B_0 by the map $\gamma' \mapsto \delta$. It is easy to see that if B is stable at level n , then so is B_1 , and that $\{\pi_n(B)\} = \{\pi_n(B_0)\} = \{\pi_n(B_1)\}\mathbb{L}^{-c}$, where π_i are truncation morphisms of $\mathcal{L}_\infty\mathcal{X}$, $L_\infty\mathcal{N}$, and $L_\infty E_1$ respectively. Therefore

$$(3.4) \quad \chi_h \mu_{E_1}(B_1) = \mu_{\mathcal{X}}(B)\mathbb{L}^c.$$

Put $\sigma := v_\infty(\delta)$. The chain of the correspondences, $\gamma \mapsto \gamma' \mapsto \delta \mapsto \sigma$, defines a map $(\mathcal{L}_\infty\mathcal{X})_{\mathcal{Y}} \rightarrow L_\infty\tilde{\mathcal{N}}$, which is the same as one in the proof of the theorem (see diagram (3.1) and compare it with the last two ones).

Shrinking $\text{Spec } R$ to an open subset, suppose that the canonical sheaf $\omega_{\text{Spec } R}$ of $\text{Spec } R$ is generated by a section e' . Consider a d -form $e = dx_1 \wedge \cdots \wedge dx_c \wedge e'$ on $\mathcal{N}$. This is stable under the G -action. If r denotes the natural morphism $\tilde{\mathcal{N}} \rightarrow \mathcal{N}$, the canonical sheaf $\omega_{\tilde{\mathcal{N}}}$ of $\tilde{\mathcal{N}}$ is generated by a d -form $\tilde{e}$ with $r^*\tilde{e} = e$. Direct computation gives $v^*\tilde{e} = t^{\sum a_i/l}(dx_1 \wedge \cdots \wedge dx_c \wedge e')$. Hence we have the following equations of subsheaves of $\Omega_{E_1[[t]]/D_\infty}^d$,

$$\begin{aligned} (t^{\sum a_i/l})v^{-1}\mathcal{J}_{\tilde{\mathcal{N}}} \cdot v^*\omega_{\tilde{\mathcal{N}}/D_\infty} &= (t^{\sum a_i/l})(v^*\Omega_{\tilde{\mathcal{N}}[[t]]/D_\infty}^d)/(tors) \\ &= (t^{\sum a_i/l})\mathcal{J}_v \cdot \Omega_{E_1[[t]]/D_\infty}^d \\ &= \mathcal{J}_v \cdot v^*\omega_{\tilde{\mathcal{N}}/D_\infty}. \end{aligned}$$

This means that $\text{ord } \mathcal{J}_{\tilde{\mathcal{N}}} \circ v_\infty - \text{ord } \mathcal{J}_v \equiv -\sum a_i/l$. From the transformation rule, we obtain that $\chi_h \mu_{\tilde{\mathcal{N}}}^{\text{Gor}}(A) = \mathbb{L}^{-\sum a_i/l} \mu_{E_1}(B_1)$, and granting (3.4), that $\chi_h \mu_{\tilde{\mathcal{N}}}^{\text{Gor}}(A) = \mathbb{L}^{s(\mathcal{Y})} \mu_{\mathcal{X}}(B)$. We have proved the assertion in this case.

As for the general case, the proof follows along almost the same lines: We take the fiber product $\mathcal{N} \times_{\mathcal{Y}} V$ for an atlas $V \rightarrow \mathcal{Y}$ to linearize the canonical automorphism. Define B_V , $B_{V,1}$ and $\tilde{B}_V$ in the evident fashion. By the argument for the preceding case and a common argument, after replacing B , we obtain that $B_{V,1}$ and $\tilde{B}_V$ are stable at level n and a morphism $v_n : \pi_n B_{V,1} \rightarrow \pi_n \tilde{B}_V$ is a trivial affine space bundle of the expected relative dimension. Here we have used Lemma 3.19 instead of the transformation rule itself. The natural morphism $\pi_{n+1} B_V \rightarrow \pi_n B_{V,1}$ is an affine space bundle of relative dimension $d - c$ which is trivial Zariski locally on V . (Recall (3.2) and (3.3). This bundle results from the truncation $L_{n+1}(\text{Spec } R) \rightarrow L_n(\text{Spec } R)$ and the identity of $(r_i)_{0 \leq i \leq n}$.) Hence $\pi_{n+1} B_V \rightarrow \pi_n \tilde{B}_V$ is also a Zariski locally trivial affine space bundle of the expected relative dimension. By the same argument as the proof of Lemma 3.11, we can conclude that

$\pi_{n+1}B \rightarrow \pi_n\tilde{B}$ is an analytically locally trivial fibration of a quotient of an affine space. The lemma follows from Lemma 3.14. $\square$

4. GENERAL RESULTS ON DELIGNE-MUMFORD STACKS

In this section, we give some general results on Deligne-Mumford stacks which we need in the preceding section. There are good references for stacks today (for example [DM69], [Vis89], [Góm01] and [LMB00]).

We fix a base scheme S .

4.1. Deligne-Mumford stacks. A *stack* is a category fibered in groupoids over (Sch/S) such that every Isom functor is a sheaf and every descent datum is effective. A morphism of stacks $\mathcal{X} \rightarrow \mathcal{Y}$ is *representable*⁴ if for any $U \in (\text{Sch}/S)$ and any morphism $U \rightarrow \mathcal{Y}$, the fiber product $U \times_{\mathcal{Y}} \mathcal{X}$ is represented by a scheme.

Definition 4.1. Let $\mathbf{P}$ be a property of morphisms $f : Y \rightarrow X$ of S -schemes, stable under base change and local in the étale topology on X (for example: surjective, proper *etc*). We say that a representable morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ of stacks has property $\mathbf{P}$ if for every S -scheme U and every morphism $U \rightarrow \mathcal{X}$, the projection $U \times_{\mathcal{X}} \mathcal{Y} \rightarrow U$ has property $\mathbf{P}$.

Definition 4.2. A (*separated*) *Deligne-Mumford stack* is a stack $\mathcal{X}$ which satisfies the following:

- (1) the diagonal $\Delta : \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is representable and finite,
- (2) there exists a scheme M and a morphism $M \rightarrow \mathcal{X}$ (necessarily representable after (1)), which is étale and surjective.

A scheme M in (2) is called an *atlas* of $\mathcal{X}$. A Deligne-Mumford stack $\mathcal{X}$ is of *finite type* if there is an atlas of finite type.

Definition 4.3. Let $\mathbf{P}$ be a property of morphisms $f : Y \rightarrow X$ of S -schemes, stable under *étale* base change and local in the étale topology on X (for example: birational, being an open immersion with dense image *etc*). We say that a representable morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ of stacks has property $\mathbf{P}$ if for every scheme U and every étale morphism $U \rightarrow \mathcal{X}$, the projection $U \times_{\mathcal{X}} \mathcal{Y} \rightarrow U$ has property $\mathbf{P}$.

Definition 4.4. Let $\mathbf{Q}$ be a property of schemes local in the étale topology (for example: reduced, smooth, normal, locally integral *etc*). Let $\mathcal{X}$ be a Deligne-Mumford stack. We say that $\mathcal{X}$ has property $\mathbf{Q}$ if an atlas of $\mathcal{X}$ has property $\mathbf{Q}$.

Definition 4.5. A (not necessarily representable) morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ of Deligne-Mumford stacks of finite type is *proper* if there is a S -scheme Z and a proper surjective morphism $g : Z \rightarrow \mathcal{Y}$ such that $f \circ g$ is (necessarily representable and) proper.

⁴In [LMB00], this is called *schématique*.

A Deligne-Mumford stack $\mathcal{X}$ of finite type is *complete* if it is proper over S .

Although our condition appears weaker than one of [DM69, Def. 4.11], they are actually equivalent by Chow's lemma (see [DM69, Def. 4.12], [LMB00, Th. 16.6], [Vis89, Prop. 2.6]).

Example 4.6. Let Z be a S -scheme and G a finite group acting on X . The *quotient stack* $[Z/G]$ is defined as follows: an object over $U \in (\text{Sch}/S)$ is a G -torsor $P \rightarrow U$ with a G -equivariant morphism $P \rightarrow Z$, and a morphism over $U' \rightarrow U$ is a cartesian diagram

$$\begin{array}{ccc} P' & \longrightarrow & P \\ \downarrow & \square & \downarrow \\ U' & \longrightarrow & U \end{array}$$

which is compatible with the G -equivariant morphisms $P' \rightarrow Z$ and $P \rightarrow Z$. It is a Deligne-Mumford stack with a canonical atlas $Z \rightarrow [Z/G]$.

Here we define points of a Deligne-Mumford stack. For details, see [LMB00, Ch. 5].

Definition 4.7. Let $\mathcal{X}$ be a Deligne-Mumford stack. A *point* of $\mathcal{X}$ is a S -morphism $\text{Spec } K \rightarrow \mathcal{X}$ for a field K with a morphism $\text{Spec } K \rightarrow S$.

Let $x_i : \text{Spec } K_i \rightarrow \mathcal{X}$ ($i = 1, 2$) be points of $\mathcal{X}$. We say that x_1 and x_2 are *equivalent* if there is a field K_3 such that $K_3 \supset K_1, K_2$ and the diagram

$$\begin{array}{ccc} \text{Spec } K_3 & \longrightarrow & \text{Spec } K_2 \\ \downarrow & & \downarrow \\ \text{Spec } K_1 & \longrightarrow & \mathcal{X} \end{array}$$

commutes.

Definition 4.8. We define the *set of points* of $\mathcal{X}$, denoted $|\mathcal{X}|$, to be the set of the equivalence classes of points of $\mathcal{X}$.

The *Zariski topology* on $|\mathcal{X}|$ is defined as follows: an open subset is $|\mathcal{U}| \subset |\mathcal{X}|$ for an open substack $\mathcal{U} \subset \mathcal{X}$. There is a 1-1 correspondence between the closed subsets of $|\mathcal{X}|$ and the reduced closed substacks of $\mathcal{X}$.

We now introduce the notion of (étale) groupoid space which is equivalent to Deligne-Mumford stack. Some references are [Vis89, p. 668], [LMB00, (2.4.3), (3.4.3), Prop. 3.8, (4.3)] and [Góm01, Subsec. 2.4].

Definition 4.9. An (étale) *groupoid space* consists of the following data:

- (1) two S -schemes X_0 and X_1 ,
- (2) five morphisms: *source* and *target* $q_i : X_1 \rightarrow X_0$ ($i = 1, 2$), *origin* $\varepsilon : X_0 \rightarrow X_1$, *inverse* $\tau : X_1 \rightarrow X_1$ and *composition* $m : X_1 \times_{q_1, X_0, q_2} X_1 \rightarrow X_1$ which satisfies the following:

- (a) q_1 and q_2 are étale and $(q_1, q_2) : X_1 \rightarrow X_0 \times X_0$ is finite,
- (b) the axioms of *associativity*, *identity element* and *inverse*.

We denote this groupoid space by $X_1 \rightrightarrows X_0$.

Given a groupoid space $X_1 \rightrightarrows X_0$, we define the category fibered in groupoids $[X_1 \rightrightarrows X_0]'$ as follows: an object over $U \in (\text{Sch}/S)$ is a morphism $U \rightarrow X_0$ of S -schemes and a morphism of $a : U \rightarrow X_0$ to $b : V \rightarrow X_0$ is a pair of morphisms $f : U \rightarrow V$ and $h : V \rightarrow X_1$ such that $q_1 \circ h \circ f = a$ and $q_2 \circ h = b$. Then $[X_1 \rightrightarrows X_0]'$ is a *prestack* (see [LMB00, 3.1]).

Definition 4.10. We define the *stack associated* to a groupoid space $X_1 \rightrightarrows X_0$, denoted $[X_1 \rightrightarrows X_0]$, to be the stack associated to the prestack $[X_1 \rightrightarrows X_0]'$ ([LMB00, Lem. 3.2]).

The stack $\mathcal{X} = [X_1 \rightrightarrows X_0]$ is a Deligne-Mumford stack with a canonical atlas $X_0 \rightarrow \mathcal{X}$. We can identify the fiber product $X_0 \times_{\mathcal{X}} X_0$ with X_1 . Conversely, given a Deligne-Mumford stack $\mathcal{X}$ and an atlas $X_0 \rightarrow \mathcal{X}$, then the schemes X_0 and $X_1 = M \times_{\mathcal{X}} M$ underlies a natural groupoid space structure with $q_i = \text{pr}_i$ and $\varepsilon = \Delta_{M/\mathcal{X}}$. The associated stack $[M \times_{\mathcal{X}} M \rightrightarrows M]$ is canonically isomorphic to $\mathcal{X}$. In summary, giving a groupoid space $X_1 \rightrightarrows X_0$ is equivalent to giving a Deligne-Mumford stack $\mathcal{X}$ and an atlas $X_0 \rightarrow \mathcal{X}$.

Let $\xi : U \rightarrow \mathcal{X}$ be a morphism, which is considered as an object of $\mathcal{X}$. If ξ lifts to $\xi' : U \rightarrow X_0$, then the automorphism group of ξ is identified with the set of morphisms $\eta : U \rightarrow X_1$ with $q_1 \circ \eta = q_2 \circ \eta = \xi'$.

Definition 4.11. A *morphism* $f : (Y_1 \rightrightarrows Y_0) \rightarrow (X_1 \rightrightarrows X_0)$ of groupoid spaces is a pair of morphisms $f_i : Y_i \rightarrow X_i$ ($i = 0, 1$) which respects the groupoid space structures.

Given a morphism $f : (Y_1 \rightrightarrows Y_0) \rightarrow (X_1 \rightrightarrows X_0)$, then we have a natural morphism of prestacks $[f]' : [Y_1 \rightrightarrows Y_0]' \rightarrow [X_1 \rightrightarrows X_0]'$ and hence a natural morphism of stacks $[f] : [Y_1 \rightrightarrows Y_0] \rightarrow [X_1 \rightrightarrows X_0]$ from [LMB00, Lem. 3.2].

Conversely, consider a commutative diagram

$$\begin{array}{ccc} Y_0 & \xrightarrow{f_0} & X_0 \\ \downarrow & & \downarrow \\ \mathcal{Y} & \xrightarrow{g} & \mathcal{X} \end{array}$$

such that $\mathcal{X}, \mathcal{Y}$ are Deligne-Mumford stacks and the vertical arrows are atlases. If we define $Y_1 := Y_0 \times_{\mathcal{Y}} Y_0$ and $X_1 := X_0 \times_{\mathcal{X}} X_0$, and if $f_1 : Y_1 \rightarrow X_1$ is the natural morphism, then the pair of (f_0, f_1) determines a morphism $f : (Y_1 \rightrightarrows Y_0) \rightarrow (X_1 \rightrightarrows X_0)$ of groupoid spaces. Evidently, $[f] = g$.

Example 4.12. Let $\mathcal{X} = [Z/G]$ be a quotient stack with G finite. There is a canonical atlas $Z \rightarrow \mathcal{X}$. Then the groupoid space $Z \times_{\mathcal{X}} Z \rightrightarrows Z$ is isomorphic to the groupoid space

$$Z \times G \begin{array}{c} \xrightarrow{G\text{-action}} \\ \xrightarrow{\text{pr}_1} \end{array} Z$$

whose origin, inverse and composition are induced by the group structure of G .

Let $\mathcal{X}$ be a locally integral Deligne-Mumford stack, associated to a groupoid space $X_1 \rightrightarrows X_0$. Let X_i^{nor} be the normalization of X_i respectively. Then the lifts of the structure morphisms of $X_1 \rightrightarrows X_0$ induce a groupoid space $X_1^{\text{nor}} \rightrightarrows X_0^{\text{nor}}$.

Definition 4.13. [Vis89, Def. 1.18] We define the *normalization* $\mathcal{X}^{\text{nor}}$ of $\mathcal{X}$ to be the stack associated to $X_1^{\text{nor}} \rightrightarrows X_0^{\text{nor}}$.

It is easy to show the uniqueness and the universality of the normalization.

4.2. Quasi-coherent sheaves.

Definition 4.14. A *quasi-coherent sheaf* $\mathcal{F}$ on a Deligne-Mumford stack $\mathcal{X}$ consists of the following data:

- (1) for each étale morphism $U \rightarrow \mathcal{X}$ with a scheme U , a quasi-coherent sheaf $\mathcal{F}_U$ on U ,
- (2) for each diagram of étale morphisms

$$\begin{array}{ccc} V & \xrightarrow{\varphi} & U \\ & \searrow & \swarrow \\ & \mathcal{X} & \end{array}$$

with V, U schemes, an isomorphism $\Theta_\varphi : \mathcal{F}_V \rightarrow \varphi^* \mathcal{F}_U$ which satisfies the cocycle condition.

- Example 4.15.** (1) The *structure sheaf* $\mathcal{O}_{\mathcal{X}}$ on $\mathcal{X}$ is defined by $(\mathcal{O}_{\mathcal{X}})_U := \mathcal{O}_U$.
- (2) The *sheaf of differentials* $\Omega_{\mathcal{X}/S}$ is defined by $(\Omega_{\mathcal{X}/S})_U := \Omega_{U/S}$ and by the canonical isomorphism.
- (3) Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a morphism of Deligne-Mumford stacks. We define the *sheaf of relative differentials* $\Omega_{\mathcal{Y}/\mathcal{X}}$ of $\mathcal{Y}$ over $\mathcal{X}$ to be the unique sheaf such that for each commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{\text{étale}} & \mathcal{Y} \\ \downarrow & & \downarrow \\ U & \xrightarrow{\text{étale}} & \mathcal{X}, \end{array}$$

$(\Omega_{\mathcal{Y}/\mathcal{X}})_V := \Omega_{V/U}$. Then we have the following exact sequence

$$f^* \Omega_{\mathcal{X}/S} \rightarrow \Omega_{\mathcal{Y}/S} \rightarrow \Omega_{\mathcal{Y}/\mathcal{X}} \rightarrow 0.$$

Definition 4.16. In Definition 4.14, if every $\mathcal{F}_U$ is an $\mathcal{O}_U$ -algebra and every Θ_φ is a homomorphism of $\mathcal{O}_V$ -algebras, then we say that $\mathcal{F}$ is an $\mathcal{O}_{\mathcal{X}}$ -algebra.

As the case of schemes, to an $\mathcal{O}_{\mathcal{X}}$ -algebra $\mathcal{F}$, we can associate a representable and affine morphism $\mathcal{S}pec \mathcal{F} \rightarrow \mathcal{X}$. For details, see [LMB00, (14.2)].

4.3. Inertia stacks. In this subsection, we study the inertia stack. It is an algebro-geometric object corresponding to the *twisted sector* which was introduced by Kawasaki [Kaw78] and used by Chen and Ruan to define the orbifold cohomology [CR00].

Definition 4.17. For a Deligne-Mumford stack $\mathcal{X}$, its *inertia stack*, denoted $I(\mathcal{X})$, is the stack defined as follows: an object over $U \in (\text{Sch}/S)$ is a pair (ξ, α) where $\xi \in \text{ob } \mathcal{X}_U$ and $\alpha \in \text{Aut}(\xi)$, and a morphism $(\xi, \alpha) \rightarrow (\eta, \beta)$ is a morphism $\gamma : \xi \rightarrow \eta$ in $\mathcal{X}$ such that $\gamma \circ \alpha = \beta \circ \gamma$.

There is a natural forgetting morphism $I(\mathcal{X}) \rightarrow \mathcal{X}$. The forgetting morphism $I(\mathcal{X}) \rightarrow \mathcal{X}$ is isomorphic to

$$\text{pr}_1 : \mathcal{X} \times_{\Delta, \mathcal{X} \times \mathcal{X}, \Delta} \mathcal{X} \rightarrow \mathcal{X}.$$

Hence if $\mathcal{X}$ is complete, then so is $I(\mathcal{X})$.

The following lemma may be well-known:

Lemma 4.18. *Let Z be a scheme with an action of a finite group G . Then we have an isomorphism:*

$$I([Z/G]) \cong \coprod_{g \in \text{Conj}(G)} [Z^g/C(g)],$$

where $\text{Conj}(G)$ is a set of representatives of the conjugacy classes, Z^g is the locus of fixed points under the g -action and $C(g)$ is the centralizer of g .

Proof. Let $U \in (\text{Sch}/S)$ be a connected scheme. An object of $[Z/G]$ over U is a G -torsor $P \rightarrow U$ with a G -equivariant morphism $P \rightarrow Z$. Its automorphism α is an automorphism of a G -torsor $P \rightarrow U$ compatible with $P \rightarrow Z$. For some étale surjective $V \rightarrow U$, the fiber product $P_V := P \times_U V$ is isomorphic to $G \times V$ as G -torsors over V . Here $G \times V$ is a G -torsor for the *right action* of G . The pull back α_V is represented by the *left action* of some $g \in G$. g is determined up to conjugacy and we can assume $g \in \text{Conj}(G)$.

Let $\psi : G \times V \cong P_V \rightarrow P$ be the natural morphism. Now let us show that if $a \in C(g)$ and $b \notin C(g)$, then we have $\psi(\{a\} \times V) \cap \psi(\{b\} \times V) = \emptyset$. Let x (resp. y) be a geometric point of $\{a\} \times V$ (resp. $\{b\} \times V$) and assume $\psi(x) = \psi(y)$. Then we have

$$\psi(x) = \psi(gxg^{-1}) = \alpha\psi(x)g^{-1} = \alpha\psi(y)g^{-1} = \psi(gyg^{-1}) \neq \psi(y).$$

This is a contradiction. So P decomposes into $C(g)$ -torsors as $P \cong P' \times J$ where $P' := \psi(C(g) \times V)$ and J is a finite set. Let f_V denote the composition $f \circ \psi$. In the following diagram,

$$\begin{array}{ccc} C(g) \times V & \xrightarrow{f_V} & Z \\ \alpha_V \downarrow & \nearrow f_V & \downarrow g \\ C(g) \times V & \xrightarrow{f_V} & Z \end{array}$$

we have $g \circ f_V = f_V \circ \alpha_V$, since α_V equals the right action of g on $C(g) \times V$ and f_V is G -equivariant. We also have $f_V = f_V \circ \alpha_V$ and hence $g \circ f_V = f_V$. It implies that $f_V(C(g) \times V)$ is in Z^g and hence so is $f(P')$. Thus the $C(g)$ -torsor $P' \rightarrow V$ with $f : P' \rightarrow Z^g$ is an object of $[Z^g/C(g)]$. For a non-connected $U \in (\text{Sch}/S)$ and an object of $I([Z/G])$ over U , we can assign it an object of $\coprod_{g \in \text{Conj}(G)} [Z^g/C(g)]$, in the obvious way. We leave the rest for the reader. $\square$

Definition 4.19. A morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ of stacks is *barely faithful* if for every object ξ of $\mathcal{Y}$, the map $\text{Aut}(\xi) \rightarrow \text{Aut}(f(\xi))$ is bijective.

Clearly, all barely faithful morphisms are faithful functors. From [LMB00, Prop. 2.3 and Cor. 8.1.2], all barely faithful morphisms of Deligne-Mumford stacks are *representable* in the sense of [LMB00, Déf. 3.9]. Because all separated and quasi-finite morphisms of algebraic spaces are schématique [LMB00, Th. A.2], all barely faithful and quasi-finite morphisms of Deligne-Mumford stacks are *representable* for our definition.

Example 4.20. All immersions are barely faithful. All morphisms of schemes are barely faithful.

Lemma 4.21. *Barely faithful morphisms are stable under base change.*

Proof. Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a barely representable morphism of stacks and $a : \mathcal{X}' \rightarrow \mathcal{X}$ any morphism of stacks. An object of the fiber product $\mathcal{Y} \times_{\mathcal{X}} \mathcal{X}'$ is a triple (ξ, η, α) , where ξ is an object of $\mathcal{Y}$, η is an object of $\mathcal{X}'$ and $\alpha : f(\xi) \rightarrow a(\eta)$ is a morphism in $\mathcal{X}_U$ for some $U \in (\text{Sch}/S)$. Its automorphism is a pair of automorphisms $\varphi \in \text{Aut}(\xi)$ and $\psi \in \text{Aut}(\eta)$ with $\alpha \circ f(\varphi) = a(\psi) \circ \alpha$. Since the map $f : \text{Aut}(\xi) \rightarrow \text{Aut}(f(\xi))$ is bijective, for each ψ , there is one and only one φ with $\alpha \circ f(\varphi) = a(\psi) \circ \alpha$. We have thus proved the lemma. $\square$

Proposition 4.22. *Let $f : \mathcal{Y} \rightarrow \mathcal{X}$ be a barely faithful morphism of Deligne-Mumford stacks. Then the inertia stack $I(\mathcal{Y})$ is naturally isomorphic to the fiber product $\mathcal{Y} \times_{\mathcal{X}} I(\mathcal{X})$.*

Proof. The natural morphism $\Psi : I(\mathcal{Y}) \rightarrow \mathcal{Y} \times_{\mathcal{X}} I(\mathcal{X})$ is defined as follows: for an object ξ of $\mathcal{Y}$ and its automorphism α , the pair (ξ, α) , which is an object of $I(\mathcal{Y})$, is mapped to the triple $(\xi, (f(\xi), f(\alpha)), \text{id}_{f(\xi)})$.

We will first show Ψ is a fully faithful functor. Let ξ be an object of $\mathcal{Y}$. The automorphism group of $(\xi, (f(\xi), f(\alpha)), \text{id}_{f(\xi)})$ is a pair of automorphisms $\beta \in \text{Aut}(\xi)$ and $\nu \in \text{Aut}((f(\xi), f(\alpha))) = C(f(\alpha))$ such that $f(\beta) = \nu$. Hence Ψ is barely faithful. Let η be another object of $\mathcal{Y}$ and τ an automorphism of η . It suffices to show that if $\text{Hom}_{I(\mathcal{Y})}((\xi, \alpha), (\eta, \tau)) = \emptyset$, then $\text{Hom}_{\mathcal{Y} \times_{\mathcal{X}} I(\mathcal{X})}(\Psi(\xi, \alpha), \Psi(\eta, \tau)) = \emptyset$. Suppose that there is an element (σ, ε) of $\text{Hom}_{\mathcal{Y} \times_{\mathcal{X}} I(\mathcal{X})}(\Psi(\xi, \alpha), \Psi(\eta, \tau))$, where σ is a morphism $\xi \rightarrow \eta$ and ε

is a morphism $f(\xi) \rightarrow f(\eta)$ such that the diagram

$$\begin{array}{ccc} f(\xi) & \xrightarrow{\varepsilon} & f(\eta) \\ f(\alpha) \downarrow & & \downarrow f(\tau) \\ f(\xi) & \xrightarrow{\varepsilon} & f(\eta) \end{array}$$

is commutative and $f(\sigma) = \varepsilon$. Since f is barely faithful, the diagram

$$\begin{array}{ccc} \xi & \xrightarrow{\sigma} & \eta \\ \alpha \downarrow & & \downarrow \tau \\ \xi & \xrightarrow{\sigma} & \eta \end{array}$$

is commutative, that is, $\text{Hom}_{I(\mathcal{Y})}((\xi, \alpha), (\eta, \tau)) \neq \emptyset$.

Now, let us show Ψ is an isomorphism, that is, an equivalence of categories. Let $(\xi, (\theta, \beta), v)$ be an object of $\mathcal{Y} \times_{\mathcal{X}} I(\mathcal{X})$ where v is an isomorphism $f(\xi) \rightarrow \theta$. Then there is a natural bijection $\text{Aut}(\xi) \rightarrow \text{Aut}(f(\xi)) \rightarrow \text{Aut}(\theta)$. Let $\alpha \in \text{Aut}(\xi)$ be an automorphism corresponding to $\beta \in \text{Aut}(\theta)$. Then we can see that $\Phi(\xi, \alpha)$ is isomorphic to $(\xi, (\theta, \beta), v)$. We have thus completed the proof. $\square$

Corollary 4.23. *If $S = \text{Spec } \mathbb{C}$ and $\mathcal{X}$ is a smooth Deligne-Mumford stack, then $I(\mathcal{X})$ is also smooth.*

Proof. From Lemma 4.26 and Lemma 4.21, there is an étale, surjective and barely faithful morphism $\coprod_i [M_i/G_i] \rightarrow \mathcal{X}$ such that each M_i is smooth and each G_i is a finite group. Then the assertion follows from Lemma 4.18 and Proposition 4.22. $\square$

4.4. Coarse moduli space.

Definition 4.24. Let $\mathcal{X}$ be a Deligne-Mumford stack. The *coarse moduli space* of $\mathcal{X}$ is an algebraic space X with a morphism $\mathcal{X} \rightarrow X$ such that:

- (1) for any algebraically closed field Ω with a morphism $\text{Spec } \Omega \rightarrow S$, $\mathcal{X}(\Omega) \rightarrow X(\Omega)$ is a bijection,
- (2) for any algebraic space Y , any morphism $\mathcal{X} \rightarrow Y$ uniquely factors as $\mathcal{X} \rightarrow X \rightarrow Y$.

Keel and Mori proved that the coarse moduli space always exists [KM97, Cor. 1.3].

Example 4.25. Let Z be an algebraic space and G a finite group acting on Z . Then the coarse moduli space of the quotient stack $[Z/G]$ is the quotient algebraic space Z/G .

The following lemma is well-known.

Lemma 4.26. *e.g. [AV02, Lem. 2.2.3] Let $\mathcal{X}$ be a Deligne-Mumford stack and X its coarse moduli space. Then there is an étale covering $(X_i \rightarrow X)_i$*

such that $\mathcal{X} \times_X X_i$ is isomorphic to a quotient stack $[Z_i/G_i]$ with a scheme Z_i and a finite group G_i . Hence the canonical morphism $\mathcal{X} \rightarrow X$ is proper.

Now we assume $S = \text{Spec } \mathbb{C}$.

Definition 4.27. Let X be a variety. We say that X has *quotient singularities* if there is an étale covering $(U_i/G_i \rightarrow X)_i$ with a smooth variety U_i and a finite group G_i .

Lemma 4.26 shows that for a variety X , X has quotient singularities if it is the coarse moduli space of some smooth Deligne-Mumford stack. In fact, ‘only if’ also holds (Lemma 4.29).

Let $\mathcal{X}$ be a smooth Deligne-Mumford stack and $x : \text{Spec } \mathbb{C} \rightarrow \mathcal{X}$ a closed point. Then $\text{Aut}(x)$ acts on the tangent space $T_x \mathcal{X}$.

Definition 4.28. We say that $\alpha \in \text{Aut}(x)$ is a *reflection* if the subspace of the α -fixed points $(T_x \mathcal{X})^\alpha$ is of codimension 1.

Lemma 4.29. Let X be a k -variety with quotient singularities. Then there is a smooth Deligne-Mumford stack $\mathcal{X}$ without reflections such that the automorphism group of general geometric points is trivial and X is the coarse moduli space of $\mathcal{X}$.

Proof. We will give only a sketch. There is a finite set of pairs $(V_i \rightarrow X, G_i)_i$ such that:

- (1) V_i is a smooth variety,
- (2) G_i is a finite group acting effectively on V_i without reflections,
- (3) $V_i \rightarrow X$ is a morphism étale in codimension 1 which factors as $V_i \rightarrow V_i/G_i \rightarrow X$ with $V_i/G_i \rightarrow X$ étale.

Let V_{ij} be the normalization of $V_i \times_X V_j$. Then the natural morphisms $V_{ij} \rightarrow V_i$ and $V_{ij} \rightarrow V_j$ are étale in codimension 1. From the purity of branch locus, they are actually étale, and hence V_{ij} is smooth. The diagonal $\Delta : V_i \rightarrow V_i \times_X V_i$ factors through $\Delta' : V_i \rightarrow V_{ii}$. Then, with the suitable multiplication morphism, the diagram

$$\coprod V_{i,j} \xrightleftharpoons{\Delta'} \coprod V_i$$

has the structure of groupoid space. We set $\mathcal{X}$ to be the associated stack. Clearly $\mathcal{X}$ has no reflections. The canonical morphism $\mathcal{X} \rightarrow X$ makes X the coarse moduli space of $\mathcal{X}$ (see [Gil84, Prop. 9.2]). Any geometric point x of $\mathcal{X}$ has a lift $\tilde{x} : \text{Spec } \Omega \rightarrow V_i$ with Ω algebraically closed field. The automorphism group of x is identified with $\{y : \text{Spec } \Omega \rightarrow V_{ii} \mid p_1 \circ y = p_2 \circ y = \tilde{x}\}$. If y is over the smooth locus of X , then this group is trivial. $\square$

REFERENCES

- [AV02] Dan Abramovich and Angelo Vistoli. Compactifying the space of stable maps. *J. Amer. Math. Soc.*, 15:27–75 (electronic), 2002.

- [Bat99a] Victor V. Batyrev. Birational Calabi-Yau n -folds have equal Betti numbers. In *New trends in algebraic geometry (Warwick, 1996)*, pages 1–11. Cambridge Univ. Press, Cambridge, 1999.
- [Bat99b] Victor V. Batyrev. Non-Archimedean integrals and stringy Euler numbers of log-terminal pairs. *J. Eur. Math. Soc. (JEMS)*, 1(1):5–33, 1999.
- [CR00] Weimin Chen and Yongbin Ruan. A new cohomology theory for orbifold. preprint, [math.AG/0004129](#), 2000.
- [Cra99] Alastair Craw. An introduction to motivic integration. preprint, [math.AG/9911179](#), 1999.
- [CR88] Charles W. Curtis and Irving Reiner. *Representation theory of finite groups and associative algebras*. John Wiley & Sons Inc., New York, 1988.
- [Dan78] Vladimir I. Danilov. The geometry of toric varieties. *Russian Math. Surveys*, 33:97–154, 1978.
- [DHVW] L. Dixon, J. A. Harvey, C. Vafa, and E. Witten. Strings on orbifolds. I, II. *Nuclear Phys. B*. 261(4):678–686, 1985. 274(2):285–314, 1986.
- [DL99] Jan Denef and François Loeser. Germs of arcs on singular algebraic varieties and motivic integration. *Invent. Math.*, 135(1):201–232, 1999.
- [DL01] Jan Denef and François Loeser. Geometry on arc spaces of algebraic varieties. In *European Congress of Mathematics, Vol. I (Barcelona, 2000)*, pages 327–348. Birkhäuser, 2001.
- [DL02] Jan Denef and François Loeser. Motivic integration, quotient singularities and the McKay correspondence. *Compositio Math.*, 131:267–290, 2002.
- [DM69] Pierre Deligne and David Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, (36):75–109, 1969.
- [EH99] David Eisenbud and Joe Harris. *The geometry of schemes*. Springer-Verlag, New York, 1999.
- [Eis95] David Eisenbud. *Commutative algebra with a view toward algebraic geometry*. Springer-Verlag, New York, 1995.
- [Gil84] Henri Gillet. Intersection theory on algebraic stacks and Q -varieties. *J. Pure Appl. Algebra*, 34(2-3):193–240, 1984.
- [Góm01] Tomás L. Gómez. Algebraic stacks. *Proc. Indian Acad. Sci. Math. Sci.*, 111(1):1–31, 2001.
- [Gre61] Marvin J. Greenberg. Schemata over local rings. *Ann. of Math. (2)*, 73:624–648, 1961.
- [EGA] Alexandre Grothendieck. *Éléments de géométrie algébrique*. Number 8, 11, 17, 20, 24, 28, 32. Inst. Hautes Études Sci. Publ. Math., Paris. written with the collaboration of Jean Dieudonné.
- [Kaw78] Tetsuro Kawasaki. The signature theorem for V -manifolds. *Topology*, 17(1):75–83, 1978.
- [KM97] Seán Keel and Shigefumi Mori. Quotients by groupoids. *Ann. of Math. (2)*, 145(1):193–213, 1997.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki. Introduction to the minimal model problem. In *Algebraic geometry, Sendai, 1985*, pages 283–360. North-Holland, Amsterdam, 1987.
- [Knu71] Donald Knutson. *Algebraic spaces*. Springer-Verlag, Berlin, 1971. Lecture Notes in Mathematics, Vol. 203.
- [Kon95] Maxim Kontsevich. Lecture at orsay. 1995.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*. Springer-Verlag, Berlin, 2000.

- [Loo02] Eduard Looijenga. Motivic measures. *Séminaire Bourbaki, Vol. 1999/2000, Astérisque*, 276:267–297, 2002.
- [Rua00] Yongbin Ruan. Stringy geometry and topology of orbifolds. preprint, **math.AG/0011149**, 2000.
- [Sai89] Morihiko Saito. Introduction to mixed Hodge modules. *Astérisque*, 179-180:145–162, 1989.
- [Sai90] Morihiko Saito. Mixed Hodge modules. *Publ. Res. Inst. Math. Sci.*, 26(2):221–333, 1990.
- [Vis89] Angelo Vistoli. Intersection theory on algebraic stacks and on their moduli spaces. *Invent. Math.*, 97(3):613–670, 1989.
- [Wat74] Keiichi Watanabe. Certain invariant subrings are Gorenstein. I, II. *Osaka J. Math.*, 11:1–8 and 379–388, 1974.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO,
TOKYO, 153-8914, JAPAN

E-mail address: `t-yasuda@ms.u-tokyo.ac.jp`